

Kirjallisuutta (kirjaan Pysy liikkuvana)

Kirjallisuusviitteitä on runsaasti. Osaan niistä on viitattu suoraan, osaa on käytetty muodostamaan laajaa ja tasapainoista kuvaa kustakin aiheesta tuoreen, korkeatasoisissa lähteissä julkaistun tieteellisen tiedon perusteella. Viitteiden kirjaamisella on pyritty helpottamaan kiinnostuneiden lukijoiden mahdollisuuksia tarkistaa, tarkentaa ja laajentaa eri aihepiirejä koskevaa tietoa ja käsityksiä.

Ravinto

Aro A, Mutanen M, Uusitupa M (toim.). Ravitsemustiede, 4.-5.painos. Kustannus Oy Duodecim, Helsinki 2015.

Autier P, Boniol M, Pizot C, Mullie P. Vitamin D statu and ill health: a systematic review. *Lancet Diab Endocrinol.* 2014;2:76-89.

Backx EMP, Tieland M, Borgonjen-van den Berg K.J, Claessen P, Loon LJC van, Groot CPGM de. Protein intake and lean body mass preservation during energy intake restriction in overweight older adults. *Int J Obes.* 2016;40:299-304.

Bischoff-Ferrari HA, Dawson-Hughes B, Orav EJ ym. Monthly high-dose vitamin D treatment for the prevention of functional decline: a randomized clinical trial. *JAMA Intern Med.* 2016;176:175-83.

Bolland MJ, Leung W, Tai V ym. Calcium intake and risk of fracture: systematic review. *BMJ.* 2015;351:h4580.

Chung M, Tang AM, Fu Z, Wang DD, Newberry SJ. Calcium intake and cardiovascular disease risk: An updated systematic review and meta-analysis. *Ann Intern Med.* 2016;165:856-866.

Del Gobbo LC, Imamura F, Aslibekyan S, ym. ω -3 Polyunsaturated fatty acid biomarkers and coronary heart disease. Pooling Project of 19 Cohort Studies. JAMA Intern Med. 2016;176:1155-1166.

Dietary Guidelines Advisory Committee. Scientific Report of the 2015 Dietary Guidelines Advisory Committee. 2015.

<http://www.health.gov/dietaryguidelines/2015-scientific-report/>. Viitattu 24.5.2017.

Ding M, Satija A, Bhupathiraju SN ym. Association of coffee consumption with total and cause-specific mortality in 3 large prospective cohorts. Circulation. 2015;132:2305-15.

Fineli – elintarvikkeiden koostumustietopankki. Terveysten ja hyvinvoinnin laitos. <https://fineli.fi/fineli/fi/index>

Finger D, Goltz FR, Umpierre D, Meyer E, Rosa LH, Schneider CD. Effects of protein supplementation in older adults undergoing resistance training: a systematic review and meta-analysis. Sports Med. 2015 ;45:245-55.

Fogelholm M, Valsta L. Ovatko transrasvat uhka kansanterveydelle?. Duodecim. 2016;132:1304-6.

Guallar E, Starnes S, Mulrow C, Appel LJ, Miller ER. Enough is enough: stop wasting money on vitamin and mineral supplements. Ann Intern Med. 2013;159:850-851-851.

Guessous I. Role of vitamin D deficiency in extraskeletal complications: predictor of health outcome or marker of health status? BioMed Res Int. 2015, Article ID 563403.

Hansen KE, Johnson E, Chambers KR ym. Treatment of vitamin D insufficiency in postmenopausal women. A randomized clinical trial. JAMA Intern Med. 2015;175:1612-1621.

He FJ, Li J, Macgregor GA. Efficacy of longer-term modest salt reduction on blood pressure. *Cochrane Database Syst Rev.* 2013;4:CD004937.

Hill TR, Aspray TJ. The role of vitamin D in maintaining bone health in older people. *Ther Adv Musculoskelet Dis.* 2017;9:89-95.

Hooper L, Martin N, Abdelhamid A, Davey Smith G. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database Syst Rev.* 2015;6:CD011737.

Khaw KT, Stewart AW, Waayer D ym. Effect of monthly high-dose vitamin D supplementation on falls and non-vertebral fractures: secondary and post-hoc outcomes from the randomised, double-blind, placebo-controlled ViDA trial. *Lancet Diabetes Endocrinol.* 2017;5:438-447.

Kyrö C, Tjønneland A. Whole grains and public health. *BMJ.* 2016;353:i3046.

Noble N, Paul C, Turon H, Oldmeadow C. Which modifiable health risk behaviours are related? A systematic review of the clustering of smoking, nutrition, alcohol and physical activity ('SNAP') health risk factors. *Prev Med.* 2015;81:16-41.

Pasiakos SM, McLellan TM, Lieberman HR. The effects of protein supplements on muscle mass, strength, and aerobic and anaerobic power in healthy adults: a systematic review. *Sports Med.* 2015;45:111-31.

Scragg R, Stewart AW, Waayer D ym. Effect of monthly high-dose vitamin D supplementation on cardiovascular disease in the Vitamin D Assessment Study : A randomized clinical trial. *JAMA Cardiol.* 2017;2:608-616.

Smith R. Are some diets “mass murder”. *BMJ.* 2014;349:g7654.

Suominen MH, Jyväkorpi SK, Pitkälä KH ym. Nutritional guidelines for older people in Finland. *J Nutr Health Aging.* 2014;18:861-7.

Tai V, Leung W, Grey A, Reid IR, Bolland M. Calcium intake and bone mineral density: systematic review and meta-analysis. *BMJ*. 2015;351:h4183.

Terveyttä ruoasta. Suomalaiset ravitsemussuositukset 2014. Valtion ravitsemusneuvottelukunta 2014.

http://www.ravitsemusneuvottelukunta.fi/files/images/vrn/2014/ravitsemus-suositukset_2014_fi_web.pdf. Viitattu 15.9.2017.

Zong G, Gao A, Hu FB, Sun Q. Whole grain intake and mortality from all causes, cardiovascular disease, and cancer: a meta-analysis of prospective cohort studies. *Circulation*. 2016;133:2370-2380.

U.S. Department of Health and Human Services and U.S. Department of Agriculture. 2015 – 2020 Dietary Guidelines for Americans. 8th Edition. December 2015. <http://health.gov/dietaryguidelines/2015/guidelines/>. Viitattu 12.3.2017.

Liikunta ja liikuntatottumukset

Liikunta

Ainsworth BE, Haskell WL, Herrmann SD ym. 2011 compendium of physical activities: a second update of codes and MET values. *Med Sci Sports Exerc*. 2011;43:1575-81.

Barnett I, van Sluijs EMF, Ogilvie D. Physical activity and transitioning to retirement: a systematic review. *Am J Prev Med*. 2012;43:329-36.

Bauman AE, Reis RS, Sallis JF, Wells JC ym. Correlates of physical activity: why are some people physically active and others not? *Lancet*. 2012;380(9838):31-33.

Bottorf JL, Seaton CL, Johnson ST ym. An updated review of interventions that include promotion of physical activity for adult men. *Sports Med*. 2015;45:775-800.

- Brawner CA, Churilla JR, Keteyian SJ. Prevalence of physical activity is lower among individuals with chronic disease. *Med Sci Sports Exerc.* 2016;48:1062-7.
- Dai S, Wang F, Morrison H. Predictors of decreased physical activity level over time among adults: a longitudinal study. *Am J Prev Med.* 2014;47:123-30.
- Das P, Horton R. Physical activity—time to take it seriously and regularly. *Lancet* 2016; 388(10051):1254-5.
- Dontje ML, Krijnen WP, de Greef MH ym. Effect of diagnosis with a chronic disease on physical activity behavior in middle-aged women. *Prev Med.* 2016;83:56-62.
- Engberg E, Alen M, Kukkonen-Harjula K, Peltonen JE, Tikkanen HO, Pekkarinen H. Life events and change in leisure time physical activity: a systematic review. *Sports Med.* 2012;42:433-47.
- French DP, Olander EK, Chisholm A, Mc Sharry J. Which behaviour change techniques are most effective at increasing older adults' self-efficacy and physical activity behaviour? A systematic review. *Ann Behav Med.* 2014;48:225-34.
- Füzéki E, Engeroff T, Banzer W. Health benefits of light-intensity physical activity: A systematic review of accelerometer data of the National Health and Nutrition Examination Survey (NHANES). *Sports Med.* 2017;47:1769-1793.
- Golubic R, Martin KR, Ekelund U ym. Levels of physical activity among a nationally representative sample of people in early old age: results of objective and self-reported assessments. *Int J Behav Nutr Phys Act.* 2014;11:58.
- Hallman DM, Birk Jørgensen M, Holtermann A. On the health paradox of occupational and leisure-time physical activity using objective

measurements: Effects on autonomic imbalance. PLoS One. 2017;12:e0177042.

Hamer M, Kivimaki M, Steptoe A. Longitudinal patterns in physical activity and sedentary behaviour from mid-life to early old age: a substudy of the Whitehall II cohort. J Epidemiol Community Health. 2012;66:1110-5.

Hansen BH, Holme I, Anderssen SA, Kolle E. Patterns of objectively measured physical activity in normal weight, overweight, and obese individuals (20-85 years): a cross-sectional study. PLoS One. 2013;8(1):e53044.

Husu P, Suni J, Vähä-Ypyä H. ym. Suomalaisten aikuisten kiihtyvyyksmittarilla mitattu fyysinen aktiivisuus ja liikkumattomuus Suomen lääkäril. 2014;69:1860-1866.

Jackson SE, Steptoe A, Wardle J. The influence of partner's behavior on health behavior change: the English Longitudinal Study of Ageing. JAMA Intern Med. 2015;175:385-92.

Jefferis BJ, Sartini C, Lee IM ym. Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. BMC Public Health. 2014;14:382.

Jefferis BJ, Sartini C, Ash S ym. Trajectories of objectively measured physical activity in free-living older men. Med Sci Sports Exerc. 2015;47:343-9.

Joutsenniemi K, Härkänen T, Pankakoski M ym. Confidence in the future, health-related behaviour and psychological distress: results from a web-based cross-sectional study of 101 257 Finns. BMJ Open. 2013;3(6). pii: e002397.

Marques A, Sarmiento H, Martins J, Saboga Nunes L. Prevalence of physical activity in European adults - compliance with the World Health Organization's physical activity guidelines. Prev Med. 2015;81:333-8.

Olander EK, Fletcher H, Williams S, Atkinson L, Turner A, French DP. What are the most effective techniques in changing obese individuals' physical activity self-efficacy and behaviour: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2013;10:29.

Pedersen BK, Saltin B. Exercise as medicine – evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015;(Suppl. 3)25:1–72.

Pedisic Z. Measurement issues and poor adjustments for physical activity and sleep undermine the sedentary behaviour research – the focus should shift to the balance between sleep, sedentary behaviour, standing and activity. *Kinesiology* 2014;46:135-146.

Portugeijs E. Käsitys omista kyvyistä ja ympäristön mahdollisuuksista määrittää ikäihmisen liikkumista ja elinpiiriä. *Liikunta & Tiede.* 2014;51:31-4.

Saarela H, Mäki-Opas T, Silventoinen K, Borodulin K. Sosiodemografiset erot suomalaisten hyötyliikunnassa. *Liikunta & Tiede.* 2015;52:55-63.

Schwaneberg T, Weymar F, Ulbricht S, Dörr M, Hoffmann W, van den Berg N. Relationship between objectively measured intensity of physical activity and self-reported enjoyment of physical activity. *Prev Med Rep.* 2017;7:162-168.

Sparling PB, Howard BJ, Dunstan DW, Owen N. Recommendations for physical activity in older adults. *BMJ.* 2015;350:h100.

Stephan AJ, Strobl R, Müller M ym. A high level of household physical activity compensates for lack of leisure time physical activity with regard to deficit accumulation: Results from the KORA-Age study. *Prev Med.* 2016;86:64-9.

Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Med.* 2014;44:81-121.

Vuori IM, Lavie CJ, Blair SN. Physical activity promotion in the health care system. *Mayo Clin Proc.* 2013;88:1446-61.

Warburton DE, Bredin SS. Reflections on physical activity and health: what should we recommend? *Can J Cardiol.* 2016;32:495-504.

Welk GJ, Kim Y. Context of physical activity in a representative sample of adults. *Med Sci Sports Exerc.* 2015;47:2102-2110.

Liikkumattomuus

Benatti FB, Ried-Larsen M. The effects of breaking up prolonged sitting time: a review of experimental studies. *Med Sci Sport Exerc.* 2015;47:2053-61.

Biswas A, Oh PI, Faulkner GE ym. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med.* 2015;162:123-32.

Dempsey PC, Howard BJ, Lynch BM, Owen N, Dunstan DW. Associations of television viewing time with adults' well-being and vitality. *Prev Med.* 2014;69:69-74.

Gibbs BB, Hergenroeder AL, Katzmarzyk PT, Lee IM, Jakicic JM. Definition, measurement, and health risks associated with sedentary behavior. *Med Sci Sports Exerc.* 2015;47:1295-300.

Healy GN, Winkler EA, Owen N, Anuradha S, Dunstan DW. Replacing sitting time with standing or stepping: associations with cardio-metabolic risk biomarkers. *Eur Heart J.* 2015;36:2643-9.

Helajärvi H, Lindholm H, Vasankari T, Heinonen OJ. Vähäisen liikkumisen terveyshaitat. *Duodecim.* 2015;131:1713-8.

Helajärvi H. Sedentary behaviour and health with special reference to obesity and fatty liver in early midlife.

<https://www.doria.fi/handle/10024/120088>. Viitattu 20.5.2017.

Husu P, Aittasalo M, Kukkonen-Harjula K. Jaloittele välillä! – perusteluja ja ratkaisuja istumisen ja muun paikallaanolon vähentämiseen. *Liikunta & Tiede*. 2016;53:18-23.

Manini TM, Carr LJ, King AC, Marshall S, Robinson TN, Rejeski WJ. Interventions to reduce sedentary behavior. *Med Sci Sports Exerc*. 2015;47:1306-10.

Mansoubi M, Pearson N, Biddle SJ, Clemes S. The relationship between sedentary behaviour and physical activity in adults: a systematic review. *Prev Med*. 2014;69:28-35.

Prince SA, Saunders TJ, Gresty K, Reid RD. A comparison of the effectiveness of physical activity and sedentary behaviour interventions in reducing sedentary time in adults: a systematic review and meta-analysis of controlled trials. *Obes Rev*. 2014;15:905-19.

Shrestha N, Kukkonen-Harjula KT, Verbeek JH, Ijaz S, Hermans V, Bhaumik S. Workplace interventions for reducing sitting at work. *Cochrane Database Syst Rev*. 2016;3:CD010912.

Stenholm S, Pulakka A, Kawachi I ym. Changes in physical activity during transition to retirement: a cohort study. *Int J Behav Nutr Phys Act*. 2016;13:51.

Tew GA, Posso MC, Arundel CE, McDaid CM. Systematic review: height-adjustable workstations to reduce sedentary behaviour in office-based workers. *Occup Med*. 2015;65:357-66.

Thyfault JP, Du M, Kraus WE, Levine JA, Booth FW. Physiology of sedentary behavior and its relationship to health outcomes. *Med Sci Sports Exerc*. 2015;47:1301-5.

Liikunnan riskit ja haitat

Chugh SS, Weiss JB. Sudden cardiac death in the older athlete. *J Am Coll Cardiol.* 2015;65:493-502.

Fitchett DH, Hegele RA, Verma S. Statin intolerance. *Circulation.* 2015;131:e389-91.

Hagel BE, Pless IB, Goulet C, Platt RW, Robitaille Y. Effectiveness of helmets in skiers and snowboarders: case-control and case crossover study. *BMJ.* 2005;330:281.

Kallinen M, Markku A. Aging, physical activity and sports injuries. An overview of common sports injuries in the elderly. *Sports Med.* 1995;20:41-52.

Kim JH, Malhotra R, Chiampas G ym. Cardiac arrest during long-distance running races. *N Engl J Med.* 2012;366:130-40.

Leifer ES, Mikus CR, Karavirta L ym Adverse cardiovascular response to aerobic exercise training: is this a concern? *Med Sci Sports Exerc.* 2016;48:20-5.

Liikunta. Käypä hoito -suositus. 13.01.2016. Suomalaisen Lääkäriseura Duodecimin ja Käypä hoito -johtoryhmän asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50075> .

Viitattu 16.4.2017.

Marijon E, Uy-Evanado A, Reinier K ym. Sudden cardiac arrest during sports activity in middle age. *Circulation.* 2015;131:1384-91.

Näveri H, Vuori I. Äkkikuoleman vaara, s. 548 – 566. Kirjassa Vuori I, Taimela S, Kujala U., toim. Liikuntalääketiede. Kustannus Oy Duodecim 2005, Helsinki.

Parkkari J. Liikuntatapaturmat, s. 567 – 579. Kirjassa Vuori I, Taimela S, Kujala U., toim. Liikuntalääketiede. Kustannus Oy Duodecim 2005 Helsinki.

Parkkari J. Liikunnan turvallisuus, s. 236-46. Kirjassa Fogelholm M, Vuori I, Vasankari T., toim. Terveysliikunta. Kustannus Oy Duodecim 2011 Helsinki.

Persaud N, Coleman E, Zwolakowski D, Lauwers B, Cass D. Nonuse of bicycle helmets and risk of fatal head injury: a proportional mortality, case-control study. CMAJ. 2012;184:E921-3.

Riebe D, Franklin BA, Thompson PD ym. Updating ACSM's recommendations for exercise preparticipation health screening. Med Sci Sports Exerc. 2015;47:2473-9.

Seron P, Lanas F, Pardo Hernandez H, Bonfill Cosp X. Exercise for people with high cardiovascular risk. Cochrane Database Syst Rev. 2014;8:CD009387.

Smith M, Matheson GO, Meeuwisse WH. Injuries in cross-country skiing: a critical appraisal of the literature. Sports Med. 1996;21:239-50.

Virtanen K. Pyöräilijän vammat. Duodecim 2016;132:1352-6.

Whang W, Manson JE, Hu F ym. Physical exertion, exercise, and sudden cardiac death in women. JAMA. 2006;295:1399-403.

Uni ja nukkumistottumukset

Burke TM, Markwald RR, McHill AW Ym. Effects of caffeine on the human circadian clock in vivo and in vitro. Sci Transl Med. 2015;7(305):305ra146.

Clark I, Landolt HP. Coffee, caffeine, and sleep: A systematic review of epidemiological studies and randomized controlled trials. Sleep Med Rev. 2016 Jan 30. pii: S1087-0792(16)00015-0.

Costello RB, Lentino CV, Boyd CC, O'Connell ML, Crawford CC, Sprengel ML, Deuster PA. The effectiveness of melatonin for promoting

healthy sleep: a rapid evidence assessment of the literature. *Nutr J*. 2014;13:106.

Dzierzewski JM, Buman MP, Giacobbi PR Jr ym. Exercise and sleep in community-dwelling older adults: evidence for a reciprocal relationship. *J Sleep Res*. 2014;23:61-8.

Garcia AN, Salloum IM. Polysomnographic sleep disturbances in nicotine, caffeine, alcohol, cocaine, opioid, and cannabis use: A focused review. *Am J Addict*. 2015;24:590-8.

Haario P. Sleep, health behaviours and weight among ageing employees: a follow-up study. Väitöskirja, 2015.

<https://helda.helsinki.fi/bitstream/handle/10138/152633/sleephea.pdf?sequence=1>. Viitattu 20.1.2017.

Halonen JI, Lallukka T, Pentti J ym. Change in job strain as a predictor of change in insomnia symptoms: analyzing observational data as a non-randomized pseudo-trial. *Sleep*. 2017;40.

Irish LA, Kline CE, Gunn HE, Buysse DJ, Hall MH. The role of sleep hygiene in promoting public health: A review of empirical evidence. *Sleep Med Rev*. 2015;22:23-36.

Irwin MR, Opp MR. Sleep health: reciprocal regulation of sleep and innate immunity. *Neuropsychopharmacology*. 2017;42:129-155.

Järnefelt H. Psykologiset hoitomuodot tehoavat unettomuuteen. *Suomen Lääkäril*. 2017;72:776-81.

Kelley GA, Kelley KS. Exercise and sleep: a systematic review of previous meta-analyses. *J Evid Based Med*. 2017;10:26-36.

Partinen M. Aikaerorasitus (jet lag). *Terveyskirjasto, Duodecim*. http://www.terveyskirjasto.fi/kotisivut/tk.koti?p_artikkeli=dlk01012&p_teos=dlk&p_osio=&p_selaus=23461. Viitattu 14.10.2017

Partinen M. Vireys, väsymys ja suorituskyky. Terveyskirjasto, Duodecim. http://www.terveyskirjasto.fi/kotisivut/tk.koti?p_artikkeli=dlk01007&p_teos=dlk&p_osio=&p_selaus=23461. Viitattu 14.10.2017.

Partonen T. Uni-valverytmin (unirytmien) häiriöt. Terveyskirjasto, Duodecim. http://www.terveyskirjasto.fi/terveyskirjasto/tk.koti?p_artikkeli=dlk00535. Viitattu 13.10.2017

Patel SR, Hayes AL, Blackwell T ym. The association between sleep patterns and obesity in older adults. *Int J Obes.* 2014;38:1159-64.

Pihl S, Aronen A-M. Unentaidot - Löydä uni ilman lääkkeitä. Kustannus Oy Duodecim, 3. uudistettu painos 2015, Helsinki.

Roguin Maor N, Alperin M, Shturman E ym. Effect of magnesium oxide supplementation on nocturnal leg cramps: a randomized clinical trial. *JAMA Intern Med.* 2017;177:617-623.

Spöndly-Nees S, Åsenlöf P, Lindberg E. High or increasing levels of physical activity protect women from future insomnia. *Sleep Med.* 2017;32:22-27.

Tsunoda K, Kitano N, Kai Y, Uchida K, Kuchiki T, Okura T, Nagamatsu T. Prospective study of physical activity and sleep in middle-aged and older adults. *Am J Prev Med.* 2015;48:662-73.

Unettomuus. Käypä hoito –suositus. 24.08.2017. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Unitutkimusseura ry:n asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50067>. Viitattu 13.10.2017.

Xiao Q, Keadle SK, Hollenbeck AR, Matthews CE. Sleep duration and total and cause-specific mortality in a large US cohort: interrelationships with physical activity, sedentary behavior, and body mass index. *Am J Epidemiol.* 2014;180:997-1006.

Watson NF, Badr MS, Belenky G ym. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep*. 2015;38:843-4.

Wigren H-K, Stenberg T. Kuinka nukkuminen elvyttää aivojamme? *Duodecim*. 2015;131:151-6.

Wunsch K, Kasten N, Fuchs R. The effect of physical activity on sleep quality, well-being, and affect in academic stress periods. *Nat Sci Sleep*. 2017;9:117-126.

Ylikoski A, Partinen M. Neurodegeneraatio ja uni. *Suomen Lääkäril*. 2017;72:281-5

Alkoholin käyttö

Aalto M, Jula A, Keinänen-Kiukaanniemi S ym. Alkoholinkäytön suitsiminen vähentää kohonneen verenpaineen haittoja. *Suomen Lääkäril*. 2015;70:2177-9.

Alkoholiongelman hoito. Käypä hoito -suositus. 4.11.2015. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Päihdelääketieteen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50028>.

Viitattu 3.10.2017.

Bell S, Daskalopoulou M, Rapsomaniki E ym. Association between clinically recorded alcohol consumption and initial presentation of 12 cardiovascular diseases: population based cohort study using linked health records. *BMJ*. 2017;356:j909.

Cao Y, Willett WC, Rimm EB, Stampfer MJ, Giovannucci EL. Light to moderate intake of alcohol, drinking patterns, and risk of cancer: results from two prospective US cohort studies. *BMJ*. 2015;351:h4238.

Castren S, Pakkanen A-M, Alho H. Alkoholien ongelmakäytön tunnistaminen ja hoito. *Suomen Lääkäril.* 2015;70:1497-1501.

Ejsing LK, Becker U, Tolstrup JS, Flensburg –Madsen T. Physical activity and risk of alcohol use disorders: results from a prospective cohort study. *Alcohol Alcohol.* 2015;50:206-12.

Hallgren M, Vancampfort D, Schuch F, Lundin A, Stubbs B. More reasons to move: exercise in the treatment of alcohol use disorders. *Front Psychiatry.* 2017;8:160.

Halonen JI, Stenholm S, Pulakka A ym. Trajectories of risky drinking around the time of statutory retirement: a longitudinal latent class analysis. *Addiction.* 2017;112:1163-1170.

Haseeb S, Alexander B, Baranchuk A. Wine and cardiovascular health: a comprehensive review. *Circulation.* 2017;136:1434-1448.

Kadlecova P, Andel R, Mikulik R, Handling EP, Pedersen NL. Alcohol consumption at midlife and risk of stroke during 43 years of follow-up. *Stroke.* 2015;46:627-33.

Knott C, Bell S, Britton A. Alcohol consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of more than 1.9 million individuals from 38 observational studies. *Diabetes Care.* 2015;38:1804-12.

Knott CS, Coombs N, Stamatakis E, Biddulph JP. All-cause mortality and the case for age specific alcohol consumption guidelines: pooled analyses of up to 10 population based cohorts. *BMJ.* 2015;350:h384.

Langballe EM, Ask H, Holmen J ym. Alcohol consumption and risk of dementia up to 27 years later in a large, population-based sample: the HUNT study, Norway. *Eur J Epidemiol.* 2015;30:1049-56.

Lillsunde P. Päihteet ja liikenneturvallisuus. *Suomen Lääkäril.* 2015;70:2615-9.

Lisha NE, Sussman S, Leventhal AM. Physical activity and alcohol use disorders. *Am J Drug Alcohol Abuse*. 2013;39:115-20.

Mostofsky E, Chahal HS, Mukamal KJ, Rimm EB, Mittleman MA. Alcohol and immediate risk of cardiovascular events. A systematic review and dose-response meta-analysis. *Circulation*. 2016;133:979-87.

Mäkelä P, Österberg E. Alkoholinkulutus, juomatavat ja alkoholipolitiikka. *Suomen Lääkäril*. 2016;71:409-15.

Perreault K, Bauman A, Johnson N, Britton A, Rangul V, Stamatakis E. Does physical activity moderate the association between alcohol drinking and all-cause, cancer and cardiovascular diseases mortality? A pooled analysis of eight British population cohorts. *Br J Sports Med*. 2017;51:651-657.

Piazza-Gardner AK, Barry AE. Examining physical activity levels and alcohol consumption: are people who drink more active? *Am J Health Prom*. 2012;26:e95-e104.

Simojoki K. Ikäihmisten alkoholinkäytön riskit, tunnistaminen ja hoito. *Suomen Lääkäril*. 2016;71:403-8.

Stockwell T, Zhao J, Panwar S, Roemer A, Naimi T, Chikritzhs T. Do “moderate” drinkers have reduced mortality risk? A systematic review and meta-analysis of alcohol consumption and all-cause mortality. *J Stud Alcohol Drugs* 2016;77:185-98.

Strandberg T, Simojoki K. Alkoholien ongelmakäyttö yli 75-vuotiailla. *Suomen Lääkäril*. 2017;72:2523-7.

Topiwala A, Allan CL, Valkanova V ym. Moderate alcohol consumption as risk factor for adverse brain outcomes and cognitive decline: longitudinal cohort study. *BMJ*. 2017;357:j2353.

Zhang C, Qin YY, Jiang H ym. Alcohol intake and risk of stroke: a dose-response meta-analysis of prospective studies. *Int J Cardiol.* 2014;174:669-77.

Zheng YL, Lian F, Shi Q ym. Alcohol intake and associated risk of major cardiovascular outcomes in women compared with men: a systematic review and meta-analysis of prospective observational studies. *BMC Public Health.* 2015;15:773.

Seppä K, Alho H, Kiianmaa K: *Alkoholiriippuvuus.* Kustannus Oy Duodecim 2013, Helsinki.

Stenberg J-H, Service H, Saiho S, Pihlaja S, Koivisto E-M, Holi M, Joffer G: *Irti alkoholista.* Kustannus Oy Duodecim 2014, Helsinki.

Tupakointi

Abrantes AM, Farris SG, Minami H, Strong DR, Riebe D, Brown RA. Acute effects of aerobic exercise on affect and smoking craving in the weeks before and after a cessation attempt. *Nicotine Tob Res.* 2017 May 15.

Agahi N, Shaw BA. Smoking trajectories from midlife to old age and the development of non-life-threatening health problems: a 34-year prospective cohort study. *Prev Med.* 2013;57:107-12.

Auer R, Vittinghoff E, Kiefe C ym. Change in physical activity after smoking cessation: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Addiction.* 2014;109:11712-83.

Boogaard S, Heymans MW, de Vet HC ym. Predictors of persistent neuropathic pain--a systematic Review. *Pain Physician.* 2015;18:433-57.

Bucholz EM, Beckman AL, Kiefe CI, Krumholz HM. Smoking status and life expectancy after myocardial infarction in the elderly. *Heart.* 2016;102:133-9.

Carter BD, Abnet CC, Feskanich D ym. Smoking and mortality – beyond established causes. *New Engl J Med.* 2015;372:631-40.

Choi NG, DiNitto DM. Role of new diagnosis, social isolation, and depression in older adults' smoking cessation. *Gerontologist.* 2015;55:793-801.

Conway TL, Cronan TA. Smoking, exercise, and physical fitness. *Prev Med.* 1992;21:723-34.

Cooper KH, Gey GO, Bottenberg RA. Effects of cigarette smoking on endurance performance. *JAMA* 1988;203:189-92.

Dallongeville J, Marécaux N, Fruchart JC, Amouyel P. Cigarette smoking is associated with unhealthy patterns of nutrient intake: a meta-analysis. *J Nutr.* 1998;128:1450-7.

Ditre JW, Brandon TH, Zale EL, Meagher MM. Pain, nicotine, and smoking: research findings and mechanistic considerations. *Psychol Bull.* 2011;137:1065-93.

Ditre JW, Heckman BW, Zale EL, Kosiba JD, Maisto SA. Acute analgesic effects of nicotine and tobacco in humans: a meta-analysis. *Pain.* 2016;157:1373-81.

Epstein KA, Viscoli CM, Spence JD ym. Smoking cessation and outcome after ischemic stroke or TIA. *Neurology.* 2017;89:1723-1729.

Farley AC, Hajek P, Lycett D, Aveyard P. Interventions for preventing weight gain after smoking cessation. *Cochrane Database Syst Rev.* 2012;1:CD006219.

Ferreira PH, Beckenkamp P, Maher CG, Hopper JL, Ferreira ML. Nature or nurture in low back pain? Results of a systematic review of studies based on twin samples. *Eur J Pain.* 2013;17:957-71.

Furlanetto KC, Mantoani LC, Bisca G ym. Reduction of physical activity in daily life and its determinants in smokers without airflow obstruction. *Respirology*. 2014;19:369-75.

Haasova M, Warren FC, Ussher M ym. The acute effects of physical activity on cigarette cravings: systematic review and meta-analysis with individual participant data. *Addiction*. 2013;108:26-37.

Haasova M, Warren FC, Ussher M ym. The acute effects of physical activity on cigarette cravings: exploration of potential moderators, mediators and physical activity attributes using individual participant data (IPD) meta-analyses. *Psychopharmacology*. 2014;231:1267-75.

Hajek P, Stead LF, West R, Jarvis M, Hartmann.-Boyce J, Lancaster T. Relapse prevention interventions for smoking cessation. *Cochrane Database Syst Rev*. 2013;8:CD003999.

Hoyt GL. Cigarette smoking: nicotine, carbon monoxide, and the physiological effects on exercise responses. *Sport Science Rev*. 2013;22:5-24.

Kabat GC, Heo M, Allison M ym. Smoking habits and body weight over the adult lifespan in postmenopausal women. *Am J Prev Med*. 2017;52:e77-e84.

Kaczynski AT, Manske SR, Mannell RC, Grewal K. Smoking and physical activity: a systematic review. *Am J Health Behav*. 2008;32:93-110.

Kaufman A, Augustson EM, Patrick H. Unraveling the relationship between smoking and weight: the role of sedentary behavior. *J Obesity*. 2012;2012, Article ID 735465.

Mons U, Muezzinler A, Gellert C ym. Impact of smoking and smoking cessation on cardiovascular events and mortality among older adults: meta-analysis of individual participant data from prospective cohort studies of the CHANCES consortium. *BMJ* 2015;350:h1551.

- Müezzinler A, Mons U, Gellert C ym. Smoking and all-cause mortality in older adults: results from the CHANCES consortium. *Am J Prev Med*. 2015;49:e53-63.
- Park EW, Tudiver FG, Campbell T. Enhancing partner support to improve smoking cessation. *Cochrane Database Syst Rev*. 2012;7:CD002928.
- Pan A, Wang Y, Talaei M, Hu FB. Relation of smoking with total mortality and cardiovascular events among patients with diabetes mellitus: A meta-analysis and systematic review. *Circulation*. 2015;132:1795-804.
- Pan A, Wang Y, Talaei M, Hu FB, Wu T. Relation of active, passive, and quitting smoking with incident type 2 diabetes: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol*. 2015;3:958-67.
- Parsons AC, Shraim M, Inglis J, Aveyard P, Hajek P. Interventions for preventing weight gain after smoking cessation. *Cochrane Database Syst Rev*. 2009;1:CD006219.
- Pujades-Rodriguez M, George J, Shah AD ym. Heterogeneous associations between smoking and a wide range of initial presentations of cardiovascular disease in 1 937360 people in England: lifetime risks and implications for risk prediction. *Int J Epidemiol*. 2015;44:129-41.
- Rautalahti M. Sähkösavukkeesta nikotiinin sähköinen annostelulaite. *Duodecim* 2017;133:229-30.
- Salonen J, Mäkitaro R, Harju T, Kaarteenaho R. Sähkösavukkeen vaikutukset terveyteen ja käyttö tupakasta vieroituksessa. *Suomen Lääkäril*. 2016;71:479-485.
- Sandvik L, Erikssen G, Thaulow E. Long term effects of smoking on physical fitness and lung function: a longitudinal study of 1393 middle aged Norwegian men for seven years. *BMJ*. 1995;311(7007):715-8.

Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between smoking and low back pain: a meta-analysis. *Am J Med.* 2010;123:87.e7-35.

Siu AL; U.S. Preventive Services Task Force. Behavioral and pharmacotherapy interventions for tobacco smoking cessation in adults, including pregnant women: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med.* 2015;163:622-34.

Sun K, Liu J, Ning G. Active smoking and risk of metabolic syndrome: a meta-analysis of prospective studies. *PLoS One.* 2012;7:e47791.

Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ.* 2014;348:g1151.

Treviño LA, Baker L, McIntosh S ym. Physical activity as a coping strategy for smoking cessation in mid-life and older adults. *Addict Behav.* 2014;39:885-8.

Tupakkatilasto 2014.

https://www.google.fi/?gws_rd=ssl#q=tupakkatilasto+2014. Viitattu 20.9.2017.

Zhong G, Wang Y, Zhang Y, Guo JJ, Zhao Y. Smoking is associated with an increased risk of dementia: a meta-analysis of prospective cohort studies with investigation of potential effect modifiers. *PLoS One.* 2015 ;10:e0118333.

Underner M, Perriot J, Peiffer G, Meurice JC. [The efficacy of physical activity as an aid to smoking cessation]. *Rev Mal Respir.* 2015;32:1016-33.

Ussher MH, Taylor AH, Faulkner GE. Exercise interventions for smoking cessation. *Cochrane Database Syst Rev.* 2014;8:CD002295.

Veldheer S, Yingst J, Zhu J, Foulds J. Ten-year weight gain in smokers who quit, smokers who continued smoking and never smokers in the United States, NHANES 2003-2012. *Int J Obes.* 2015;39:1727-32.

Wang D, Wang Y, Wang Y, Li R, Zhou C. Impact of physical exercise on substance use disorders: a meta-analysis. *PLoS One.* 2014;9:e110728.

Elintapoja koskevat suositukset ja niiden omakohtainen käyttö

Suominen MH, Jyväkorpi SK, Pitkälä KH ym. Nutritional guidelines for older people in Finland. *J Nutr Health Aging.* 2014;18:861-7.

Terveyttä ruoasta. Suomalaiset ravitsemussuositukset 2014. Valtion ravitsemusneuvottelukunta 2014.

http://www.ravitsemusneuvottelukunta.fi/files/images/vrn/2014/ravitsemus_suosituks_2014_fi_web.pdf. Viitattu 15.9.2017.

U.S. Department of Health and Human Services and U.S. Department of Agriculture. 2015 – 2020 Dietary Guidelines for Americans. 8th Edition. December 2015. <http://health.gov/dietaryguidelines/2015/guidelines/>. Viitattu 12.3.2017.

Liikunta. Käypä hoito -suositus. 13.01.2016. Suomalaisen Lääkäriseura Duodecimin ja Käypä hoito -johtoryhmän asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituks/suositus?id=hoi50075> .

Viitattu 16.4.2017.

WHO. Global recommendations on physical activity for health.

www.who.int/dietphysicalactivity/publications/97892. Viitattu 17.4.2017.

Unettomuus. Käypä hoito –suositus. 24.08.2017. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Unitutkimusseura ry:n asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituks/suositus?id=hoi50067>.

Viitattu 13.10.2017.

Alkoholiongelmaisen hoito. Käypä hoito -suositus. 4.11.2015. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Päihdelääketieteen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50028>.

Viitattu 3.10.2017.

Tupakkariippuvuus ja tupakasta vieroitus. 19.01.2012. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Yleislääketieteen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi40020>.

Viitattu 12.10.2017.

Alaselkäkipu

Alaselkäkipu. Käypä hoito -suositus.10.09.2014. Suomalaisen Lääkäriseura Duodecimin ja Suomen Fysiatriryhdistyksen asettama työryhmä. <http://www.kaypahoito.fi/web/kh/suosituksset/> .

[suositus?id=hoi20001](http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi20001). Viitattu 15.9.2017.

Amlani NM, Munir F. Does physical activity have an impact on sickness absence? A review. Sports Med. 2014;47:887-907.

Belavý DL, Albracht K, Bruggemann GP, Vergroesen PP, van Dieën JH. Can exercise positively influence the intervertebral disc? Sports Med. 2016;46:473-85.

Bohman T, Alfredsson L, Hallqvist J, Vingård E, Skillgate E. The influence of self-reported leisure time physical activity and the body mass index on recovery from persistent back pain among men and women: a population-based cohort study. BMC Public Health. 2013;13:385.

Bohman T, Alfredsson L, Jensen I, Hallqvist J, Vingård E, Skillgate E. Does a healthy lifestyle behaviour influence the prognosis of low back pain among men and women in a general population? A population-based cohort study. BMJ Open. 2014;4:e005713.

Brumitt J, Matheson JW, Meira EP. Core stabilization exercise prescription, part 2: a systematic review of motor control and general (global) exercise rehabilitation approaches for patients with low back pain. *Sports Health*. 2013;5:510-3.

Brämberg EB, Bergström G, Jensen I, Hagberg J, Kwak L. Effects of yoga, strength training and advice on back pain: a randomized controlled trial. *BMC Musculoskelet Disord*. 2017;18:132.

Carey TS, Freburger JK. Exercise and the prevention of low back pain: ready for implementation. *JAMA Intern Med*. 2016;176:208-9.

Cherkin DC, Sherman KJ, Balderson BH ym. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA*. 2016;315:1240-9.

Chou R, Deyo R, Friedly J ym. Noninvasive treatments for low back pain. *Comparative Effectiveness Reviews*, No. 169. Agency for Healthcare Research and Quality (US), 2016..

Coenen P, Parry S, Willenberg L ym. Associations of prolonged standing with musculoskeletal symptoms-A systematic review of laboratory studies. *Gait Posture*. 2017;58:310-318.

Delitto A, George SZ, Van Dillen LR ym. Low back pain. *J Orthop Sports Phys Ther*. 2012;42:A1-57

Fritz JM, Magel JS, McFadden M ym. Early physical therapy vs usual care in patients with recent-onset low back pain: a randomized clinical trial. *JAMA*. 2015;314:1459-67.

Furlan AD, Giraldo M, Baskwill A, Irvin E, Imamura M. Massage for low-back pain. *Cochrane Database Syst Rev*. 2015;9:CD001929.

Gordon R, Bloxham S. A systematic review of the effects of exercise and physical activity on non-specific chronic low back pain. *Healthcare*. 2016;4. pii: E22.

Grunnesjö MI, Bogefeldt JP, Blomberg SI, Strender LE, Svärdsudd KF. A randomized controlled trial of the effects of muscle stretching, manual therapy and steroid injections in addition to 'stay active' care on health-related quality of life in acute or subacute low back pain. *Clin Rehabil*. 2011;25:999-1010.

Hagen KB, Dagfinrud H, Moe RH ym. Exercise therapy for bone and muscle health: an overview of systematic reviews. *BMC Medicine* 2012;10:167.

Hoffmann TC, Maher CG, Briffa T ym. Prescribing exercise interventions for patients with chronic conditions. *CMAJ*. 2016;188:510-8.

Karppinen J, Arokoski J, Paananen M, Lausmaa M, Takala E-P, Pohjolainen T. Yksilöityjä menetelmiä epäspesifisen alaselkävun diagnostiikkaan ja hoitoon. *Duodecim*. 2016;132:1427-31

Kuijper PP, Verbeek JH, Visser B ym. An evidence-based multidisciplinary practice guideline to reduce the workload due to lifting for preventing work-related low back pain. *Ann Occup Environ Med*. 2014;26:16.

Kumar S, Beaton K, Hughes T. The effectiveness of massage therapy for the treatment of nonspecific low back pain: a systematic review of systematic reviews. *Int J Gen Med*. 2013;6:733-41.

Lawford BJ, Walters J, Ferrar K. Does walking improve disability status, function, or quality of life in adults with chronic low back pain? A systematic review. *Clin Rehabil*. 2016;30:523-36.

Lehtola V, Luomajoki H, Leinonen V, Gibbons S, Airaksinen O. Subclassification based specific movement control exercises are superior to general exercise in sub-acute low back pain when both are combined with

manual therapy: A randomized controlled trial. BMC Musculoskelet Disord. 2016;17:135.

Liikunta. Käypä-hoito suositus. 13.01.2016. Suomalaisen Lääkäriseura Duodecimin ja Käypä hoito-johtoryhmän asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50075>.

Viitattu 12.9.2017.

Lin HT, Hung WC, Hung JL, Wu PS, Liaw LJ, Chang JH. Effects of pilates on patients with chronic non-specific low back pain: a systematic review. J Phys Ther Sci. 2016;28:2961-2969.

Liu L, Skinner M, McDonough S, Mabire L, Baxter GD. Acupuncture for low back pain: an overview of systematic reviews. Evid Based Complement Alternat Med. 2015;2015:328196.

Macedo LG, Bostick GP, Maher CG. Exercise for prevention of recurrences of nonspecific low back pain. Phys Ther. 2013;93:1587-91.

Macedo LG, Saragiotto BT, Yamato TP, Costa LO, Menezes Costa LC, Ostelo RW, Maher CG. Motor control exercise for acute non-specific low back pain. Cochrane Database Syst Rev. 2016;2:CD012085.

Maher C, Underwood M, Buchbinder R. Non-specific low back pain. Lancet. 2017;389(10070):736-747.

McGorry RW, Hsiang SM, Snook SH, Clancy EA, Young SL. Meteorological conditions and self-report of low back pain. Spine 1998;23:2096-102.

Morone NE, Greco CM, Moore CG ym. A mind-body program for older adults with chronic low back pain: a randomized clinical trial. JAMA Intern Med. 2016;176:329-37.

Nelson NL. Kinesio taping for chronic low back pain: A systematic review. J Bodyw Mov Ther. 2016;20:672-81.

Paanalahti K, Holm LW, Nordin M, Asker M, Lyander J, Skillgate E. Adverse events after manual therapy among patients seeking care for neck and/or back pain: a randomized controlled trial. *BMC Musculoskeletal Disord.* 2014;15:77.

Paige NM, Miake-Lye IM, Booth MS ym. Association of spinal manipulative therapy with clinical benefit and harm for acute low back pain: systematic review and meta-analysis. *JAMA.* 2017;317:1451-1460.

Parreira P, Heymans MW, van Tulder M ym. Back schools for chronic non-specific low back pain. *Cochrane Database Syst Rev.* 2017;8:CD011674.

Poquet N, Lin CW, Heymans MW ym. Back schools for acute and subacute non-specific low-back pain. *Cochrane Database Syst Rev.* 2016;4:CD008325.

Qaseem A, Wilt TJ, McLean RM, Forciea MA; Clinical Guidelines Committee of the American College of Physicians. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2017;166:514-530.

Saragiotto BT, Maher CG, Yamato TP, Costa LO, Menezes Costa LC, Ostelo RW, Macedo LG. Motor control exercise for chronic non-specific low-back pain. *Cochrane Database Syst Rev.* 2016;(1):CD012004.

SBU. Preventiva insatser vid akut smärta från rygg och nacke – Effekter av fysisk träning, manuell behandling och beteendepåverkande åtgärder. Stockholm: Statens beredning för medicinsk och social utvärdering (SBU). 2016. <http://www.sbu.se/sv/publikationer/SBU-utvarderar/preventiva-insatser-vid-akut-smarta-fran-rygg-och-nacke--effekter-av-fysisk-traning-manuell-behandling-och-beteendepaverkande-atgarder/>. Viitattu 12.5.2017.

Shiri R, Lallukka T, Karppinen J, Viikari-Juntura E. Obesity as a risk factor for sciatica: a meta-analysis. *Am J Epidemiol.* 2014;179:929-37.

Shiri R, Coggon D, Falah-Hassani K. Exercise for the prevention of low back pain: systematic review and meta-analysis of controlled trials. *Am J Epidemiol.* 2017 Oct 19.

Shiri R, Coggon D, Falah-Hassani K. Exercise for the prevention of low back and pelvic girdle pain in pregnancy: A meta-analysis of randomized controlled trials. *Eur J Pain.* 2018;22:19-27.

Shiri R, Falah-Hassani K. Does leisure time physical activity protect against low back pain? Systematic review and meta-analysis of 36 prospective cohort studies. *Br J Sports Med.* 2017;51:1410-1418.

Steffens D, Ferreira ML, Latimer J ym. What triggers an episode of acute low back pain? A case-crossover study. *Arthritis Care Res.* 2015;67:403-10.

Steffens D, Maher CG, Pereira LS. Prevention of low back pain: a systematic review and meta-analysis. *JAMA Intern Med.* 2016;176:199-208.

Stilwell P, Harman K. Contemporary biopsychosocial exercise prescription for chronic low back pain: questioning core stability programs and considering context. *J Can Chiropr Assoc.* 2017;61:6-17.

Taulaniemi A, Kuusinen L, Tokola K, Kankaanpää M, Suni JH. Bio-psychosocial factors are associated with pain intensity, physical functioning, and ability to work in female healthcare personnel with recurrent low back pain. *J Rehabil Med.* 2017;49:667-676.

van Middelkoop M, Rubinstein SM, Kuijpers T ym. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J.* 2011;20:19-39.

Wang XQ, Zheng JJ, Yu ZW ym. A meta-analysis of core stability exercise versus general exercise for chronic low back pain. PLoS One. 2012;7(12):e52082.

Wells C, Kolt GS, Marshall P, Hill B, Bialocerkowski A. Effectiveness of Pilates exercise in treating people with chronic low back pain: a systematic review of systematic reviews. BMC Med Res Methodol. 2013;13:7.

Wieland LS, Skoetz N, Pilkington K, Vempati R, D'Adamo CR, Berman BM. Yoga treatment for chronic non-specific low back pain. Cochrane Database Syst Rev. 2017;1:CD010671.

Yamato TP, Maher CG, Saragiotto BT ym. Pilates for low back pain. Cochrane Database Syst Rev. 2015;7:CD010265.

Fibromyalgia

Ablin J, Fitzcharles M-A, Buskila D, Shir Y, Sommer C, Häuser W. Treatment of fibromyalgia syndrome: Recommendations of recent evidence-based interdisciplinary guidelines with special emphasis on complementary and alternative therapies. Evid Based Complement Alternat Med. 2013; 2013:485272.

Bawa FL, Mercer SW, Atherton RJ ym. Does mindfulness improve outcomes in patients with chronic pain? Systematic review and meta-analysis. Br J Gen Pract. 2015;65(635):e387-400.

Bidonde J, Busch AJ, Webber SC ym. Aquatic exercise training for fibromyalgia. Cochrane Database Syst Rev. 2014;10:CD011336.

Bidonde J, Busch AJ, Schachter CL ym. Aerobic exercise training for adults with fibromyalgia. Cochrane Database Syst Rev. 2017;6:CD012700.

Busch AJ, Webber SC, Brachaniec M ym. Exercise therapy for fibromyalgia. Curr Pain Headache Rep. 2011;15:358-67.

- Busch AJ, Webber SC, Richards RS ym. Resistance exercise training for fibromyalgia. *Cochrane Database Syst Rev.* 2013;12:CD010884.
- Clauw DJ. Fibromyalgia. A clinical review. *JAMA.* 2014;311:1547-1555.
- Deare JC, Zheng Z, Xue CC ym. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev.* 2013;5:CD007070.
- Ellingson LD, Shields MR, Stegner AJ, Cook DB. Physical activity, sustained sedentary behavior, and pain modulation in women with fibromyalgia. *J Pain.* 2012;13:195-206.
- Ellingson LD, Stegner AJ, Schwabacher IJ, Koltyn KF, Cook DB. Exercise strengthens central nervous system modulation of pain in fibromyalgia. *Brain Sci.* 2016;6(1). pii: E8.
- Jones KD. Recommendations for resistance training in patients with fibromyalgia. *Arthritis Res Ther.* 2015 ;17:258.
- Kivikoski L, Hannonen P. Fibroamyalgian taustalla on kivunsäätelyn ja autonomisen hermoston herkistyminen. *Suomen Lääkäril.* 2013;68:1409-1415.
- Larsson A, Palstam A, Löfgren M ym. Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgia--a randomized controlled trial. *Arthritis Res Ther.* 2015;17:161.
- Larun L, Brurberg KG, Odgaard-Jensen J, Price JR. Exercise therapy for chronic fatigue syndrome. *Cochrane Database Syst Rev.* 2015;2:CD003200.
- Lauche R, Cramer H, Häuser W, Dobos G, Langhorst J. A systematic review and meta-analysis of qigong for the fibromyalgia syndrome. *Evid Based Complement Alternat Med.* 2013;2013:635182.
- Lauche R, Cramer H, Häuser W, Dobos G, Langhorst J. A systematic overview of reviews for complementary and alternative therapies in the

treatment of the fibromyalgia syndrome. *Evid Based Complement Alternat Med.* 2015;2015:610615.

Li YH, Wang FY, Feng CQ1, Yang XF, Sun YH. Massage therapy for fibromyalgia: a systematic review and meta-analysis of randomized controlled trials. *PLoS One.* 2014;9:e89304.

Macfarlane GJ, Kronisch C, Atzeni F ym. EULAR recommendations for management of fibromyalgia. *Ann Rheum Dis.* 2017;76:e54.

Mali M. Eväitä fibromyalgian kohtaamiseen. *Suomen Lääkäril.* 2016; 71:1371-4.

Mist SD, Firestone KA, Jones KD. Complementary and alternative exercise for fibromyalgia: a meta-analysis. *J Pain Res.* 2013;6:247-60.

Naumann J, Sadaghiani C. Therapeutic benefit of balneotherapy and hydrotherapy in the management of fibromyalgia syndrome: a qualitative systematic review and meta-analysis of randomized controlled trials. *Arthritis Res Ther.* 2014;16:R141.

Nüesch E, Häuser W, Bernardy K, Barth J, Jüni P. Comparative efficacy of pharmacological and non-pharmacological interventions in fibromyalgia syndrome: network meta-analysis. *Ann Rheum Dis.* 2013; 72:955-62.

Rahman A, Underwood M, Carnes D. Fibromyalgia. *BMJ.* 2014;348:g1224.

Salehi A, Hashemi N, Imanieh MH, Saber M. Chiropractic: is it efficient in treatment of diseases? Review of systematic reviews. *Int J Community Based Nurs Midwifery.* 2015;3:244-54.

Sharif K, Watad A, Bragazzi NL, Lichtbroun M, Amital H, Shoenfeld Y. Physical activity and autoimmune diseases: Get moving and manage the disease. *Autoimmun Rev.* 2018;17:53-72.

Sosa-Reina MD, Nunez-Nagy S, Gallego-Izquierdo T, Pecos-Martín D, Monserrat J, Álvarez-Mon M. Effectiveness of therapeutic exercise in

fibromyalgia syndrome: a systematic review and meta-analysis of randomized clinical trials. *Biomed Res Int*. 2017;2017:2356346.

Theadom A, Copley M, Smith HE, Feigin VL, McPherson K. Mind and body therapy for fibromyalgia. *Cochrane Database Syst Rev*. 2015;4:CD001980.

Jännityspäänsärky

Andersen LL, Mortensen OS, Zebis MK, Jensen RH, Poulsen OM. Effect of brief daily exercise on headache among adults -- secondary analysis of a randomized controlled trial. *Scand J Work Environ Health*. 2011 ;37:547-50.

Andersen LL, Saervoll CA, Mortensen OS ym. Effectiveness of small daily amounts of progressive resistance training for frequent neck/shoulder pain: randomised controlled trial. *Pain*. 2011;152:440-6.

Atula S. Jännityspäänsärky. http://www.terveyskirjasto.fi/terveyskirjasto/tk.koti?p_artikkeli=dlk00024. Viitattu 17.10.2017.

Baillie LE, Gabriele JM, Penzien DB. A systematic review of behavioral headache interventions with an aerobic exercise component. *Headache*. 2014;54:40-53.

Bendtsen L, Fernandez-de-la-Penas C. The role of muscles in tension-type headache. *Curr Pain Headache Rep*. 2011;15:451-8.

Coeytaux RR, Befus D. Role of acupuncture in the treatment or prevention of migraine, tension-type headache, or chronic headache disorder. *Headache* 2016;56:1238-40.

Daenen L, Varkey E, Kellmann M, Nijs J. Exercise, not to exercise, or how to exercise in patients with chronic pain. *Applying science to practice. Clin J Pain*. 2015;3:108-14.

- de Groot FM, Voogt-Bode A, Passchier J, Berger MY, Koes BW, Verhagen AP. Headache: the placebo effects in control groups in randomized controlled trials; an analysis of systemic reviews. *J Manipulative Physiol Ther.* 2011;34:297-305.
- Derry S, Wiffen PJ, Moore RA. Aspirin for acute treatment of episodic tension-type headache in adults. *Cochrane Database Syst Rev.* 2017;1:CD011888.
- Gil-Martínez A, Kindelan-Calvo P, Agudo-Carmona D ym. Therapeutic exercise as treatment for migraine and tension-type headaches: a systematic review of randomised clinical trials. *Rev Neurol.* 2013;57:433-43.
- Haque B, Rahman KM, Hoque A ym. Precipitating and relieving factors of migraine versus tension type headache. *BMC Neurol.* 2012;12:82.
- Hindiye NA, Krusz JC, Cowan RP. Does exercise make migraines worse and tension type headaches better? *Curr Pain Headache Rep.* 2013;17:380.
- Häuser W, Hansen E, Enck P. Nocebo phenomena in medicine: their relevance in everyday clinical practice. *Dtsch Arztebl Int.* 2012;109:459-65.
- Linde K, Allais G, Brinkhaus B ym. Acupuncture for the prevention of tension-type headache. *Cochrane Database Syst Rev.* 2016;4:CD007587.
- Mesa-Jiminez JA, Lozano-Lopez C, Angulo-Diaz-Parreno S, Rodriguez-Fernandez AL, De_la-Hoz-Aizpurua JL, Fernandez-de-Las-Penas C. Multimodal manual therapy versus pharmacological care for management of tension type headache: A meta-analysis of randomized trials. *Cephalalgia* 2015;35:1323-32.
- Mitsikostas DD, Mantonakis LI, Chalarakis NG. Nocebo is the enemy, not placebo. A meta-analysis of reported side effects after placebo treatment in headaches. *Cephalalgia* 2011;31:550-61.

Nicholson RA, Buse DC, Andrasik F, Lipton RB. Nonpharmacologic treatments for migraine and tension-type headache: how to choose and when to use. *Curr Treat Options Neurol.* 2011;13:28-40.

Posadzki P, Ernst E. Spinal manipulations for tension-type headaches: A systematic review of randomized controlled trials. *Comp Ther Med.* 2012;20:232-9.

Stephens G, Derry S, Moore RA. Paracetamol (acetaminophen) for acute treatment of episodic tension-type headache in adults. *Cochrane Database Syst Rev.* 2016;6:CD011889.

Varatharajan S, Ferguson B, Chrobak K ym. Are non-invasive interventions effective for the management of headaches associated with neck pain? An update of the Bone and Joint Decade Task Force on Neck Pain and Its Associated Disorders by the Ontario Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Eur Spin J.* 2016;25:1971-99.

Varkey E. On the prevention of migraine. University of Gothenburg, Göteborg 2012. ISBN: 978-91-628-8406-2.

Veys L, Derry S, Moore RA. Ketoprofen for episodic tension-type headache in adults. *Cochrane Database Syst Rev.* 2016;9:CD012190.

Winter AC, Hoffmann W, Meisinger C ym. Association between lifestyle factors and headache. *J Headache Pain.* 2011;12:147–155.

Wöber C, Wöber-Bingöl C. Triggers of migraine and tension-type headache. *Handb Clin Neurol.* 2010;97:161-72.

Jäykkyys, kankeus, liikkuvuus ja liikevajaukset

Ayala F, de Baranda Andújar PS. Effect of 3 different active stretch durations on hip flexion range of motion. *J Strength Cond Res.* 2010;24:430-6.

Behm DG, Blazevich AJ, Kay AD, McHugh M. Acute effects of muscle stretching on physical performance, range of motion, and injury incidence in healthy active individuals: a systematic review. *Appl Physiol Nutr Metab.* 2016;41:1-11.

Behm DG, Cavanaugh T, Quigley P, Reid JC, Nardi PS, Marchetti P. Acute bouts of upper and lower body static and dynamic stretching increase non-local joint range of motion. *Eur J Appl Physiol.* 2016;116:241-9.

Bleakley CM, Costello JT. Do thermal agents affect range of movement and mechanical properties in soft tissues? A systematic review. *Arch Phys Med Rehabil.* 2013;94:149-63.

Chaouachi A, Padulo J, Kasmi S, Othmen AB, Chatra M, Behm DG. Unilateral static and dynamic hamstrings stretching increases contralateral hip flexion range of motion. *Clin Physiol Funct Imaging.* 2017;37:23-29.

Cipriani DJ, Terry ME, Haines MA, Tabibnia AP, Lyssanova O. Effect of stretch frequency and sex on rate of gain and rate of loss in muscle flexibility during a hamstring stretching program: a randomized single-blind longitudinal study. *J Strength Cond Res.* 2012;26:2119-29

Covert CA, Alexander MP, Petronis JJ, Davis DS. Comparison of ballistic and static stretching on hamstring muscle length using an equal stretching dose. *J Strength Cond Res.* 2010;24:3998-3014.

Cruz-Ferreira A, Fernandes J, Laranjo L, Bernardo L, Silva A. A systematic review of the effects of Pilates method of exercise in healthy people. *Arch Phys Med Rehabil.* 2011;92:2071-81.

de Baranda PS, Ayala F. Chronic flexibility improvement after 12 week of stretching program utilizing the ACSM recommendations: hamstring flexibility. *Int J Sports Med.* 2010;31:389-96.

González-Ravé JM, Sánchez-Gómez A, Santos-García DJ. Efficacy of two different stretch training programs (passive vs. proprioceptive

neuromuscular facilitation) on shoulder and hip range of motion in older people. *J Strength Cond Res.* 2012;26:1045-51.

Harvey LA, Katalinic OM, Herbert RD, Moseley AM, Lannin NA, Schuss K. Stretch for the treatment and prevention of contractures. *Cochrane Database Syst Rev.* 2017,1:CD007455.

Herbert RD, de Noronha M, Kamper SJ. Stretching to prevent or reduce muscle soreness after exercise. *Cochrane Database Syst Rev.* 2011;7:CD004577.

Johnson AW, Mitchell UH, Meek K, Feland JB. Hamstring flexibility increases the same with 3 or 9 repetitions of stretching held for a total time of 90 s. *Phys Ther Sport.* 2014;15:101-5.

Lewis J. A systematic literature review of the relationship between stretching and athletic injury prevention. *Orthop Nurs.* 2014;33:312-20.

Nakano J, Yamabayashi C, Scott A, Reid WD. The effect of heat applied with stretch to increase range of motion: a systematic review. *Phys Ther Sport.* 2012;13:180-8.

Petrofsky JS1, Laymon M, Lee H. Effect of heat and cold on tendon flexibility and force to flex the human knee. *Med Sci Monit.* 2013;19:661-7.

Stathokostas L, Little RMD, Vandervoort AA, Paterson DH. Flexibility training and functional ability in older adults: a systematic review. *J Aging Res.* 2012;Article ID 306818, 30 pages.

Weppler CH, Magnusson SP. Increasing muscle extensibility: a matter of increasing length or modifying sensation. *Phys Ther.* 2010;90:438-449.

Kaatumiset ja tasapaino

Barker AL, Bird ML, Talevski . Effect of pilates exercise for improving balance in older adults: a systematic review with meta-analysis. *Arch Phys Med Rehabil.* 2015;96:715-23.

Barker AL, Talevski , Bohensky MA, Brand CA, Cameron PA, Morello RT. Feasibility of Pilates exercise to decrease falls risk: a pilot randomized trial in community-dwelling older people. *Clin Rehabil.* 2016;30:984-96.

Bea J, Thomson CA, Wallace RB ym. Changes in physical activity, sedentary time, and risk of falling: the Women's Health Initiative Observational Study. *Prev Med.* 2017;95:103-9.

Behm DG, Muehlbauer T, Kibele A, Granacher U. Effects of strength training using unstable surfaces on strength, power and balance performance across the lifespan: a systematic review and meta-analysis. *Sports Med.* 2015;45:1645-69.

Bouillon R. Can vitamin D prevent falls and fractures? *Lancet Diabetes Endocrinol.* 2017;5:407-409.

Buchner DM, Rillamas-Sun E, Di C, LaMonte MJ ym. Accelerometer-measured moderate to vigorous physical activity and Iicidence rates of falls in older women. *J Am Geriatr Soc.* 2017;65:2480-2487.

Bullo V, Bergamin M, Gobbo S. ym. The effects of Pilates exercise training on physical fitness and wellbeing in the elderly: a systematic review for future exercise prescription. *Prev Med.* 2015;75:1-11.

Caban-Martinez AJ, Courtney TK, Chang WR ym. Leisure-time physical activity, falls, and fall injuries in middle-aged adults. *Am J Prev Med.* 2015;49:888-901.

Cadore EL, Rodriquez-Manas L, Sinclair A, Izquierdo M. Effects of different exercise interventions on risk of falls, gait ability, and balance in physically frail older adults: a systematic review. *Rejuvenation Tes.* 2013;16:105-14.

Clemson L, Fiatarone Singh MA, Bundy A ym. Integration of balance and strength training into daily activity to reduce rate of falls in older people (The LIFE study): a randomized controlled trial. *BMJ*. 2012;345:e4547.

Del-Pino-Casade R, Obero-Gaitan E, Lomas-Vega R. The effect of tai chi on reducing the risk of falling: a systematic review and meta-analysis. *Am J Chin Med*. 2016;44:895-906.

de Olivera Fransisco C, de Almeida Fagundes A, Gorges B. Effects of Pilates method in elderly people: Systematic review of randomized controlled trials. *J Bodyw Mov Ther*. 2015;19:500-8.

Donath L, Rössler R, Faude o. Effects of virtuality training (Exergaming) compared to alternative exercise training and passive control on standing balance and functional mobility in healthy community-dwelling seniors: A meta-analytical review. *Sports Med*. 2016;46:1293-1309.

El-Khoury F, Cassou B, Charles M-A, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2013;347:f6234

El-Khoury F, Cassou B, Latouche A, Aegerter P, Charles MA, Dargent-Molina P. Effectiveness of two year balance training programme on prevention of fall induced injuries in at risk women aged 75-85 living in community: Ossébo randomised controlled trial. *BMJ*. 2015;351:h3830.

Fairhall N, Sherrington C, Clemson L, Cameron ID. Do exercise interventions designed to prevent falls affect participation in life roles? A systematic review and meta-analysis. *Age Ageing* 2011;40:666-74.

Gawler S, Skelton DA, Dinan-Young S ym. Reducing falls among older people in general practice: The ProAct65+ exercise intervention trial. *Arch Gerontol Geriatr*. 2016;67:46-54.

Gianoudis J, Bailey C, Ebeling P ym. Effects of a targeted multimodal exercise program incorporating high-speed power training on falls and

fracture risk factors in older adults: A community-based randomized controlled trial. *J Bone Miner Res.* 2014;29:182-191.

Gill TM, Pahor M, Guralnik JM ym. Effect of structured physical activity on prevention of serious fall injuries in adults aged 70-89: randomized clinical trial (LIFE Study). *BMJ.* 2016;352:i245.

Gillespie LD, Robertson MC, Gillespie WJ ym. Interventions for preventing falls in older people living in the community. *Cochrane Database Sys Rev.* 2012;9:CD 007146.

Granacher U, Muehlbauer T, Gruber M. A qualitative review of balance and strength performance in healthy older adults: impact for testing and training. *J Aging Res.* Vol. 2012, Article ID 708905.

Granacher U, Gollhofer A, Hortobágyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: a systematic review. *Sports Med.* 2013;43:627-41.

Harvey NC, Biver E, Kaufman JM. ym. The role of calcium supplementation in healthy musculoskeletal aging. *Osteoporosis Int.* 2017;28:447.

Hill KD, Hunter SW, Batcelor FA, Cavalheri V, Burton E. Individualized home-based exercise programs for older people to reduce falls and improve physical performance: A systematic review and meta-analysis. *Maturitas* 2015;82:72-84.

Hoffmann TC, Maher CG, Briffa T ym. Prescribing exercise interventions for patients with chronic conditions. *CMAJ.* 2016; 188:510-8.

Hortobagyi T, Lesinski M, Gäbler M, VanSwaeringen JM, Malatesta M, Granacher U. Effects of three types of exercise interventions on healthy old adults' gait speed: A systematic review and meta-analysis. *Sports Med.* 2015;45:1627-43.

Huang Y, Liu X. Improvement of balance control ability and flexibility in the elderly Tai Chi Chuan (TCC) practitioners: a systematic review and meta-analysis. *Arch Gerontol Geriatr.* 2015;60:233-8

Hwang H-F, Chen S-J, Lee-Hsieh J, Chien D-K, Chen C-Y, Lin M-R. Effects of home-based tai chi and lower extremity training and self-practice on falls and functional outcomes in older fallers from emergency department – a randomized controlled trial. *J Am Geriatr Soc.*2016;64:518-25.

Jefferis BJ, Merom D, Sartini C, Wannamethee SG, Ash S, Lennon LT, Iliffe S, Kendrick D, Whincup PH. Physical activity and falls in older men: the critical role of mobility limitations. *Med Sci Sports Exerc.* 2015 ;47:2119-28.

Karinkanta S. To keep fit and function. Effects of three exercise programs on multiple risk factors for falls and related fractures in home-dwelling older women. *Acta Electronica Universitatis Tamperensis* 1105. (2011). ISBN 978-951-44-8536-7 (pdf). <http://acta.uta.fi>. Viitattu 12.1.2017.

Kauranen K. Tasapaino ja fysioterapia tasapaino-ongelmissa. Kirjassa Kauranen K. Fysioterapeutin käsikirja. Sanoma Pro 2017, Helsinki.

Kemmler W, Häberle L, von Stengel S. Effects of exercise on fracture reduction in older adults: a systematic review and meta-analysis. *Osteoporos Int.* 2013;24:1937-50.

Kendrick D, Kumar A, Carpenter H ym. Exercise for reducing fear of falling in older people living in the community. *Cochrane Database Syst Rev.* 2014;11:CD009848.

Korhonen Niina. Fall-induced injuries and deaths among older Finns between 1970 and 2012. Väitöskirja. *Acta Electronica Universitatis Tamperensis*; 1486, Tampere University Press 2014. ISBN 978-951-44-9638-7, ISSN 1456-954X. <http://tampub.uta.fi>. Viitattu 14.9.2017.

Kummel J, Kramer A, Giboin L-S, Gruber M. Specificity of balance training in healthy individuals: a systematic review and meta-analysis. *Sports Med.* 2016;46:1261-71.

Leung DP, Chan CK, Tsang HW, Tsang WW, Jones AY. Tai chi as an intervention to improve balance and reduce falls in older adults: A systematic and meta-analytical review. *Altern Ther Health Med.* 2011;17:40-8.

Lesinski M, Hortobágyi T, Muehlbauer T, Gollhofer A, Granacher U. Dose-response relationships of balance training in healthy young adults: a systematic review and meta-analysis. *Sports Med.* 2015;45:557-76.

Lesinski M, Hortobágyi T, Muehlbauer T, Gollhofer A, Granacher U. Effects of balance training on balance performance in healthy older adults: a systematic review and meta-analysis. *Sports Med.* 2015;45:1721-38.

Low DC, Walsh GS, Arkestejin M. Effectiveness of exercise interventions to improve postural control in older adults: a systematic review and meta-analyses of centre of pressure measurements. *Sports Med.* 2017;47:101-112.

Mertz KJ, Lee D, Sui X, Powell KE, Blair SN. Falls among adults. The association of cardiorespiratory fitness and physical activity with walking-related falls. *Am J Prev Med.* 2010;39:15-24.

Muehlbauer T, Gollhofer A, Granacher U. Associations between measures of balance and lower-extremity muscle strength/power in healthy individuals across the lifespan: a systematic review and meta-analysis. *Sports Med.* 2015;45:1671-92

Orr R, Raymand J, Fiatarone Singh M. Efficacy of progressive resistance training on balance performance in older adults: a systematic review of randomized controlled trials. *Sport Med.* 2008;38:317-43.

Osteoporoosi. Käypä hoito -suositus. 24.4.2014. Suomalaisen Lääkäriseuran Duodecimin, Suomen Endokrinologiyhdistyksen ja Suomen

Gynekologiyhdistyksen asettama työryhmä.

<http://www.terveysportti.fi/xmedia/hoi/hoi24065.pdf>. Viitattu 15.5.2017.

Pajala S, Piirtola M, Karinkanta S ym. Kaatumisten ja kaatumisvammojen ehkäisyn fysioterapiasuositus.

http://www.terveysportti.fi/dtk/sfs/avaa?p_artikkeli=sfs00003. Viitattu 15.5.2017

Pajala S. Ikäihmisten kaatumisten ehkäisy. Terveyden ja hyvinvoinnin laitos 2012, Tampere.

Patil R. Exercise in older women: effects on falls, function, fear of falling and finances. Acta Electronica Universitatis Tamperensis 1592. ISBN 978-951-44-9918-0 (pdf). ISSN 1456-954X.

Plummer P, Zukowski LA, Giuliani C, Hall AM, Zurakowski D. Effects of physical exercise interventions on gait-related dual-task interference in older adults: a systematic review and meta-analysis. Gerontology. 2015;62:94-117.

Pope Z, Zang N, Gao Z. The effects of video games on patients' rehabilitative outcomes: A meta-analysis. Prev Med. 2017;95:38-46.

Rimland JM, Abraha I, Dell'Áquila G. ym. Effectiveness of non-pharmacological interventions to prevent falls in older people: A systematic overview. The SENATOR Project ONTOP Series. PLOS One 2016;11:e0161579.

Robertson MC, Gillespie LD. Fall prevention in community-dwelling older adults. JAMA. 2013;309:1406-7.

Sherrington C, Michaleff ZA, Fairhall N ym. Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. Br J Sports Med. 2017;51:1750-1758.

Shier V, Trieu E, Ganz DA. Implementing exercise programs to prevent falls: systematic descriptive review. Injury Epidemiology 2016;3:16.

Stubbs B, Brefka S, Denkinger MD. What works to prevent falls in community-dwelling older adults? Umbrella review of meta-analyses of randomized controlled trials. *Phys Ther.* 2015;95:1095-110.

Tricco AC, Thomas SM, Veroniki AA ym. Comparisons of interventions for preventing falls in older adults: a systematic review and meta-analysis. *JAMA.* 2017;318:1687-1699.

Turner S, Kisser R, Rogmans W. Falls among older adults in the EU-28: Key facts from the available statistics. Fact Sheet, EuroSafe 2015.

Zhao R, Feng F, Wang X. Exercise interventions and prevention of fall-related fractures in older people: a meta-analysis of randomized controlled trials. *Int J Epidemiol.* 2017;46:149-161.

Zheng YT, Cui QQ, Hong YM, Yao WG. A meta-analysis of high dose, intermittent vitamin D supplementation among older adults. *PLOS One* 2015;10:e 0115850.

Uusi-Rasi K, Patil R, Karinkanta S, Kannus P, Tokola K, Lamberg-Allardt C, Sievänen H. Exercise and vitamin D in fall prevention among older women: a randomized clinical trial. *JAMA Intern Med.* 2015;175:703-11.

Uusi-Rasi K, Patil R, Karinkanta S ym. A 2-year follow-up after a 2-year RCT with vitamin D and exercise: effects on falls, injurious falls and physical functioning among older women. *J Gerontol A Biol Sci Med Sci.* 2017;72:1239-1245.

van het Reve E, de Bruin ED. Strength-balance supplemented with computerized cognitive training to improve dual task gait and divided attention in older adults: a multicenter randomized-controlled trial. *BMC Geriatr.* 2014;14:134.

Vieira ER, Palmer RC, Chaves PHM. Prevention of falls in older people living in the community. *BMJ* 2016;353:i1419.

Vlaeyen E, Coussement J, Leysens G ym. Characteristics and effectiveness of fall prevention programs in nursing homes: a systematic review and meta-analysis of randomized controlled trials. *J Am Geriatr Soc.* 2015;63:211-21.

White AM, Tooth LR, Peeters GMEEG. Fall risk factors in mid-age women: The Australian Longitudinal Study on Women's Health. *Am J Prev Med.* 2018;54:51-63.

www.turvallinenkoti.net

Youkhana S, Dean CM, Wolff M, Sherrington C, Tiedemann A. Yoga-based exercise improves balance and mobility in people aged 60 and over: a systematic review and meta-analysis. *Age Ageing* 2016;45:21-9.

Kihti

Andres M, Sivera F, Falzon L, Buchbinder R, Carmona L. Dietary supplements for chronic gout. *Cochrane Database Syst. rev.* 2014;10:CD010156.

Chen J-H, Wen CP, Wu SB ym. Attenuating the mortality risk of high serum uric acid: the role of physical activity underused. *Ann Rheum Dis.* 2015;74:2034-42.

Dalbeth N, Merriman TR, Stamp LK. Gout. *Lancet.* 2016;388(10055):2039-2052.

Julkunen H. Kihti ja valekihti. *Duodecim* 2017;133:953-6

Li X, Meng X, Timofeeva M ym. Serum uric acid levels and multiple health outcomes: umbrella review of evidence from observational studies, randomised controlled trials, and Mendelian randomisation studies. *BMJ.* 2017;357:j2376.

Mustajoki P. Kihti. Terveyskirjasto 2016.

http://www.terveyskirjasto.fi/terveyskirjasto/tk.koti?p_artikkeli=dlk00032.

Viitattu 25.5.2017.

Stewart S, Dalbeth N, McNair P, Parmar P, Gow P, Rome K. The effect of good and poor walking shoe characteristics on plantar pressure and gait in people with gout. *Clin Biochem*. 2014;29:1158-63.

Williams PT. Effects of diet, physical activity and performance, and body weight on incident gout in ostensibly healthy, vigorously active men. *Am J Clin Nutr*. 2008;87:1480-7.

Kipu

Ambrose KR, Golightly YM. Physical exercise as non-pharmacological treatment of chronic pain: Why and when. *Best Pract Res Clin Rheumatol*. 2015;29:120-30. 23.

Booth J, Moseley GL, Schiltenswolf M, Cashin A, Davies M, Hübscher M. Exercise for chronic musculoskeletal pain: A biopsychosocial approach. *Musculoskeletal Care*. 2017;15:413-421.

Cooper MA, Kluding PM, Wright DE. Emerging relationships between exercise, sensory nerves, and neuropathic pain. *Front Neurosci*. 2016 23;10:372.

Daenen L, Varkey E, Kellmann M, Nijs J. Exercise, not to exercise, or how to exercise in patients with chronic pain? Applying science to practice. *Clin J Pain*. 2015;31:108-14.

Ferrari ML, Thuraisingam S, von Känel R, Egloff N. Expectations and effects of a single yoga session on pain perception. *Int J Yoga*. 2015;8:154-7.

Geneen LJ, Moore RA, Clarke C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews. *Cochrane Database Syst Rev.* 2017;4:CD011279.

Haanpää M. Neuropaattisen kivun hoito-opas. Artikkelin tunnus: nix00086 (000.000), © 2017 Suomalainen Lääkäriseura Duodecim.

Haanpää M, Salminen JJ. Kipu, s. 54-73. Teoksessa *Fysiatria*, Arokoski J, Alaranta H, Pohjolainen T, Salminen J, Viikari-Juntura E., toim.. Duodecim 2009,Helsinki.

Kauranen K. Fysikaaliset hoidot kivun hoidossa. Teoksessa Kauranen K. *Fysioterapeutin käsikirja*, Sanoma Pro 2017, Helsinki.

Kipu. Käypä hoito -suositus. 22.08.2017. Suomalaisen Lääkäriseuran Duodecimin, Suomen Anestesiologiyhdistyksen ja Suomen Yleislääketieteen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50103>.

Viitattu 12.10.2017.

Koho P, Orenius T, Kautiainen H, Haanpää M, Pohjolainen T, Hurri H. Association of fear of movement and leisure-time physical activity among patients with chronic pain. *J Rehabil Med.* 2011;43:794-9.

Korshøj M, Birk Jørgensen M, Lidegaard M ym. Decrease in musculoskeletal pain after 4 and 12 months of an aerobic exercise intervention: a worksite RCT among cleaners. *Scand J Public Health.* 2017:1403494817717833.

Landmark T, Romundstad PR, Borchgrevink PC, Kaasa S, Dale O. Longitudinal associations between exercise and pain in the general population--the HUNT pain study. *PLoS One.* 2013;8:e65279.

Micalos PS, Arendt-Nielsen L. Differential pain response at local and remote muscle sites following aerobic cycling exercise at mild and moderate intensity. *Springerplus.* 2016 ;5:91.

Miranda H. Ota kipu haltuun. Otava 2017, Helsinki. ISBN: 9789511292302.

Nijs J, Kosek E, Van Oosterwijck J, Meeus M. Dysfunctional endogenous analgesia during exercise in patients with chronic pain: to exercise or not to exercise? *Pain Physician*. 2012;15(3 Suppl):ES205-13.

O'Connor SR, Tully MA, Ryan B ym. Walking exercise for chronic musculoskeletal pain: systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2015;96:724-734.e3.

Sundstrup E, Jakobsen MD, Brandt M, Jay K, Aagaard P, Andersen LL. Strength training improves fatigue resistance and self-rated health in workers with chronic pain: a randomized controlled trial. *Biomed Res Int*. 2016;2016:4137918.

Lihaskato eli sarkopenia

ACSM. Position Stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc*.

2009;41:1510-1530.

Aggio DA, Sartini C, Papcosta O ym. Cross-sectional associations of objectively measured physical activity and sedentary time with sarcopenia and sarcopenic obesity in older men. *Prev Med*. 2016;91:264-272.

Beaudart C, Dawson A, Shaw SC ym. Nutrition and physical activity in the prevention and treatment of sarcopenia: systematic review. *Osteoporos Int*. 2017;28:1817-1833.

Beaudart C, McCloskey E, Bruyère O ym. Sarcopenia in daily practice: assessment and management. *BMC Geriatr*. 2016;16:170.

Beaudart C, Zaaria M, Pasleau F, Reginster JY, Bruyere O. Health outcomes of sarcopenia: A systematic review and meta-analysis. *PLOS One* 2017;12:e0169548

Bickel CS, Cross JM, Bamman MM. Exercise dosing to retain resistance training adaptations in young and older adults. *Med Sci Sports Exerc.* 2011;43:1177-87.

Bleakley C, McDonough S, Gardner E, Baxter GD, Hopkins JT, Davison GW. Cold-water immersion (cryotherapy) for preventing and treating muscle soreness after exercise. *Cochrane Database Syst Rev.* 2012;2:CD008262.

Cederholm T, Cruz-Jentoft AJ, Maggi S. Sarcopenia and fragility fractures. *Eur J Phys Rehabil Med.* 2013;49:111-7.

Counts BR, Buckner SL, Mouser JG ym. Muscle growth: To infinity and beyond? *Muscle Nerve.* 2017 ;56:1022-1030.

Cruz-Jentoft AJ, Landi F, Schneider SM ym. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWGSOP and IWGS). *Age Ageing.* 2014;43:748-59.

Davies T, Orr R, Halaki M, Hackett D. Effect of training leading to repetition failure on muscular strength: a systematic review and meta-analysis. *Sports Med.* 2016;46:487-502.

English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care.* 2010;13:34-9.

Ferrucci L, Baroni M, Ranchelli A ym. Interaction between bone and muscle in older persons with mobility limitations. *Curr Pharm Des.* 2014;20:3178-97.

Fragala MS, Kenny AM, Kuchel GA. Muscle quality in aging: a multi-dimensional approach to muscle functioning with applications for treatment. *Sports Med.* 2015;45:641-58.

Granacher U, Gollhofer A, Hortobágyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: a systematic review. *Sports Med*. 2013;43:627-41.

Holviala J, Häkkinen A, Alen M, Sallinen J, Kraemer W, Häkkinen K. Effects of prolonged and maintenance strength training on force production, walking, and balance in aging women and men. *Scand J Med Sci Sports*. 2014;24:224-33.

Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol*. 2014;2:819-29.

Korpela M, Petterson Tom, Strandberg T, Löfberg M, Kiuru-Enari S. Vanhusten lihasongelmat. *Suomen Lääkärilehti*. 2011;66:2409-16.

Kwan P. Sarcopenia, a neurogenic syndrome? *J Aging Res*. 2013;2013:791679.

Leong DP, Teo KK, Rangarajan S ym. Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet*. 2015;386(9990):266-73.

Liu CJ, Latham N. Adverse events reported in progressive resistance strength training trials in older adults: 2 sides of a coin. *Arch Phys Med Rehabil*. 2010;91:1471-3.

Loenneke JP, Rossow LM, Fahs CA, Thiebaud RS, Grant Mouser J, Bemben MG. Time-course of muscle growth, and its relationship with muscle strength in both young and older women. *Geriatr Gerontol Int*. 2017;17:2000-2007.

Lozano-Montoya I, Correa-Pérez A, Abraha I. Nonpharmacological interventions to treat physical frailty and sarcopenia in older patients: a systematic overview - the SENATOR Project ONTOP Series. *Clin Interv Aging*. 2017;12:721-740.

Manini TD, Clark BC. Dynapenia and aging: an update. *J Gerontol A Biol Sci Med Sci*. 2012;67A:28-40.

Mansi I, Frei CR, Pugh MJ, Makris U, Mortensen EM. Statins and musculoskeletal conditions, arthropathies, and injuries. *JAMA Intern Med*. 2013;173:1-10.

Montero-Fernández N, Serra-Rexach JA. Role of exercise on sarcopenia in the elderly. *Eur J Phys Rehabil Med*. 2013;49:131-43.

Morton RW, Murphy KT, McKellar SR ym. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med*. 2017 Jul 11. pii: bjsports-2017-097608.

Narici MV, Maffulli N. Sarcopenia: characteristics, mechanisms and functional significance *Br Med Bull*. 2010;95:139-59.

Robinson MM, Dasari S, Konopka AR ym. Enhanced protein translation underlies improved metabolic and physical adaptations to different exercise training modes in young and old humans. *Cell Metab*. 2017;25:581-592.

Sayer AA, Robinson SM, Patel HP, Shavlakadze T, Cooper C, Grounds MD. New horizons in the pathogenesis, diagnosis and management of sarcopenia. *Age Ageing*. 2013;42:145-50.

Sénéchal M, McGavock JM, Church TS, Lee DC, Earnest CP, Sui X, Blair SN. Cut points of muscle strength associated with metabolic syndrome in men. *Med Sci Sports Exerc*. 2014;46:1475-81.

Stewart VH, Saunders DH, Greig CA. Responsiveness of muscle size and strength to physical training in very elderly people: a systematic review. *Scand J Med Sci Sports*. 2014;24:e1-10.

Sjöblom S. Postmenopausaalisen lihaskadon (sarkopenia) yhteys osteoporoosiin, murtumiin ja kaatumisiin.

http://epublications.uef.fi/pub/urn_nbn_fi_uef-20140509/urn_nbn_fi_uef-20140509.pdf. Viitattu 12.2.2017.

Theou O, Stathokostas L, Roland KP, Jakobi JM, Patterson C, Vandervoort AA, Jones GR. The effectiveness of exercise interventions for the management of frailty: a systematic review. *J Aging Res.* 2011;2011:569194.

Verdijk LB, Snijders T, Holloway TM, van Kranenburg J, Van Loon LJC. Resistance training increases skeletal muscle capillarization in healthy older men. *Med Sci Sports Exerc.* 2016;48:2157-64.

Vincent HK, Raiser SN, Vincent KR. The aging musculoskeletal system and obesity-related considerations with exercise. *Ageing Res Rev.* 2012;11:361-73.

Weinheimer EM, Sands LP, Campbell WW. A systematic review of the separate and combined effects of energy restriction and exercise on fat-free mass in middle-aged and older adults: implications for sarcopenic obesity. *Nutr Rev.* 2010;68:375-88.

Liikapaino ja lihavuus

Ahern AL, Wheeler GM, Aveyard P ym. Extended and standard duration weight-loss programme referrals for adults in primary care (WRAP): a randomised controlled trial. *Lancet.* 2017;389(10085):2214-2225. 3.

Al Khatib HK, Harding SV, Darzi J, Pot GK. The effects of partial sleep deprivation on energy balance: a systematic review and meta-analysis. *Eur J Clin Nutr.* 2017 ;71:614-624.

Apovian CM. The obesity epidemic--understanding the disease and the treatment. *N Engl J Med.* 2016;374:177-9.

Azimi A, Charlot MG, Torp-Pedersen C ym. Moderate overweight is beneficial and severe obesity detrimental for patients with documented atherosclerotic heart disease. *Heart*. 2013;99:655-60.

Backholer K, Pasupathi K, Wong E, Hodge A, Stevenson C, Peeters A. The relationship between body mass index prior to old age and disability in old age. *Int J Obes*. 2012;36:1180-86.

Backx EM, Tieland M, Borgonjen-van den Berg KJ, Claessen PR, van Loon LJ, de Groot LC. Protein intake and lean body mass preservation during energy intake restriction in overweight older adults. *Int J Obes* . 2016;40:299-304.

Banack HR, Stokes A. The 'obesity paradox' may not be a paradox at all. *Int J Obes*. 2017;41:1162-1163.

Barone Gibbs B, Pettee Gabriel K, Carnethon MC ym. Sedentary time, physical activity, and adiposity: cross-sectional and longitudinal associations in CARDIA. *Am J Prev Med*. 2017;53:764-771.

Bell JA, Kivimaki M, Hamer M. Metabolically healthy obesity and risk of incident type 2 diabetes: a meta-analysis of prospective cohort studies. *Obes Rev*. 2014;15:504-15.

Bell JA, Sabia S, Singh-Manoux A, Hamer M, Kivimäki M. Healthy obesity and risk of accelerated functional decline and disability. *Int J Obes*. 2017;41:866-872.

Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5•24 million UK adults. *Lancet*. 2014;384:755–65.

Bray GA, Frühbeck G, Ryan DH, Wilding JP. Management of obesity. *Lancet*. 2016 387(10031):1947-56.

Britton KA, Lee IM, Wang L ym. Physical activity and the risk of becoming overweight or obese in middle-aged and older women. *Obesity*. 2012;20:1096-1103.

Brown V, Moodie M, Mantilla Herrera AM, Veerman JL, Carter R. Active transport and obesity prevention - A transportation sector obesity impact scoping review and assessment for Melbourne, Australia. *Prev Med*. 2017;96:49-66.

Casazza K, Fontaine KR, Astrup, A. ym. Myths, presumptions, and facts about obesity. *N Engl J Med*. 2013; 368:446-454.

Castro EA, Júdice PB, Silva AM, Teixeira PJ, Benito PJ. Sedentary behavior and compensatory mechanisms in response to different doses of exercise-a randomized controlled trial in overweight and obese adults. *Eur J Clin Nutr*. 2017;71:1393-1398.

Caudwell P, Gibbons C, Finlayson G, Näslund E, Blundell J. No sex differences in body weight response to exercise. *Exerc Sport Sci Rev*. 2014;42:92-101.

Charakida M, Khan T, Johnson W ym. Lifelong patterns of BMI and cardiovascular phenotype in individuals aged 60—64 years in the 1946 British birth cohort study: an epidemiological study. *Lancet Diabetes & Endocrinology*. 2014;2:648–654.

Chowdhury EA, Richardson JD, Tsintzas K, Thompson D, Betts JA. Effect of extended morning fasting upon ad libitum lunch intake and associated metabolic and hormonal responses in obese adults. *Int J Obes*. 2016;40:305-11.

Cloostermans L, Picavet HS, Bemelmans WJ, Verschuren WM, Proper KI. Psychological distress as a determinant of changes in body mass index over a period of 10 years. *Prev Med*. 2015;77:17-22.

Dale CE, Fatemifar G, Palmer TM ym. Causal associations of adiposity and body fat distribution with coronary heart disease, stroke subtypes, and

type 2 diabetes mellitus: A Mendelian randomization analysis. *Circulation*. 2017;135:2373-2388.

Dandanell S, Ritz C, Verdich E, Dela F, Helge JW. Repeated lifestyle interventions lead to progressive weight loss: A retrospective review chart study. *Scand J Public Health*. 2017;45:305-313.

Dhurandhar EJ, Kaiser KA, Dawson JA, Alcorn AS, Keating KD, Allison DB. Predicting adult weight change in the real world: a systematic review and meta-analysis accounting for compensatory changes in energy intake or expenditure. *Int J Obes*. 2015;39:1181-7.

Dhurandhar NV, Schoeller D, Brown AW. Energy balance measurement: when something is not better than nothing. *Int J Obes*. 2015;39:1109-13.

Dombrowski SU, Knittle K, Avenell A, Araújo-Soares V, Sniehotta FF. Long term maintenance of weight loss with non-surgical interventions in obese adults: systematic review and meta-analyses of randomised controlled trials. *BMJ*. 2014;348:g2646.

Donges CE, Duffield R. Effects of resistance or aerobic exercise training on total and regional body composition in sedentary overweight middle-aged adults. *Appl Physiol Nutr Metab*. 2012;37:499-509.

Donnelly JE, Herrmann SD, Lambourne K, Szabo AN, Honas JJ, Washburn RA. Does increased exercise or physical activity alter ad-libitum daily energy intake or macronutrient composition in healthy adults? A systematic review. *PLoS One*. 2014;9:e83498.

Douglas JA, King JA, Clayton DJ ym. Acute effects of exercise on appetite, ad libitum energy intake and appetite-regulatory hormones in lean and overweight/obese men and women. *Int J Obes*. 2017;41:1737-1744.

Dowd JB, Zajacova A. Long-term obesity and physical functioning in older Americans. *Int J Obes*. 2015;39:502-7.

Drenowatz C, Hand GA, Sagner M, Shook RP, Burgess S, Blair SN. The prospective association between different types of exercise and body composition. *Med Sci Sports Exerc.* 2015;47:2535-41.

Drenowatz C, Hill JO, Peters JC, Soriano-Maldonado A, Blair SN. The association of change in physical activity and body weight in the regulation of total energy expenditure. *Eur J Clin Nutr.* 2017;71:377-382.

Döring N, de Munter J, Rasmussen F. The associations between overweight, weight change and health related quality of life: Longitudinal data from the Stockholm Public Health Cohort 2002-2010. *Prev Med.* 2015;75:12-7.

Ekelund U, Kolloe E, Steene-Johannessen J ym. Objectively measured sedentary time and physical activity and associations with body weight gain: does body weight determine a decline in moderate and vigorous intensity physical activity? *Int J Obes.* 2017;41:1769-1774.

Fall T, Hägg S, Ploner A ym. Age- and sex-specific causal effects of adiposity on cardiovascular risk factors. *Diabetes.* 2015;64:1841-52.

Ferraro ZM, Patterson S, Chaput JP ym. Unhealthy weight control practices: culprits and clinical recommendations. *Clin Med Insights Endocrinol Diabetes.* 2015;17:7-11.

Flint E, Cummins S. Active commuting and obesity in mid-life: cross-sectional, observational evidence from UK Biobank. *Lancet Diabetes Endocrinol.* 2016 Mar 16. pii: S2213-8587(16)00053-X.

GBD 2015 Obesity Collaborators. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med.* 2017;377:13-27.

Greenway FL. Physiological adaptations to weight loss and factors favouring weight regain. *Int J Obes.* 2015;39:1188-96.

Heymsfield SB, Wadden TA. Mechanisms, pathophysiology, and management of obesity. *N Engl J Med.* 2017;376:254-266.

Hutfless S1, Gudzone KA, Maruthur N ym. Strategies to prevent weight gain in adults: a systematic review. *Am J Prev Med.* 2013;45:e41-51.

Jackson SE, Beeken RJ, Wardle J. Obesity, perceived weight discrimination, and psychological well-being in older adults in England. *Obesity.* 2015;23:1105-11.

Jensen MD, Ryan DH, Apovian CM ym. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol.* 2014;63:2985-3023.

Keating SE, Johnson NA, Mielke GI, Coombes JS. A systematic review and meta-analysis of interval training versus moderate-intensity continuous training on body adiposity. *Obes Rev.* 2017;18:943-964.

Kivimäki M, Kuosma E, Ferrie JE. Overweight, obesity, and risk of cardiometabolic multimorbidity: pooled analysis of individual-level data for 120 813 adults from 16 cohort studies from the USA and Europe. *Lancet Public Health.* 2017;2:e277-e285.

Kramer CK, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions? A systematic review and meta-analysis. *Ann Intern Med.* 2013;159:758-69.

Kyrgiou M, Kalliala I, Markozannes G ym. Adiposity and cancer at major anatomical sites: umbrella review of the literature. *BMJ.* 2017;356:j477.

Larsen B, Allison MA, Kang E ym. Associations of physical activity and sedentary behavior with regional fat deposition. *Med Sci Sports Exerc.* 2014;46:520-28.

Lihavuus (aikuiset). Käypä hoito -suositus. 13.09.2013. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Lihavuustutkijat ry:n asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suositukset/suositus?id=hoi24010>.

Viitattu 4.10.2017.

Luke A, Cooper RS. Physical activity does not influence obesity risk: time to clarify the public health message. *Int J Epidemiol.* 2013;42:1831-6.

Lv N, Azar KMJ, Rosas LG, Wulfovich S, Xiao L, Ma J. Behavioral lifestyle interventions for moderate and severe obesity: A systematic review. *Prev Med.* 2017;100:180-193.

Ma C, Avenell A, Bolland M ym. Effects of weight loss interventions for adults who are obese on mortality, cardiovascular disease, and cancer: systematic review and meta-analysis. *BMJ.* 2017;359:j4849.

Mabire L, Mani R, Liu L, Mulligan H, Baxter D. The influence of age, sex and body mass index on the effectiveness of brisk walking for obesity management in adults: a systematic review and meta-analysis. *J Phys Act Health.* 2017;14:389-407.

Madigan CD, Daley AJ, Lewis AL, Aveyard P, Jolly K. Is self-weighing an effective tool for weight loss: a systematic literature review and meta-analysis. *Int J Behav Nutr Phys Act.* 2015;12:104.

Magkos F, Fraterrigo G, Yoshino J. Effects of moderate and subsequent progressive weight loss on metabolic function and adipose tissue biology in humans with obesity. *Cell Metab.* 2016;23:591-601.

Malhotra A, Noakes T, Phinney S. It is time to bust the myth of physical inactivity and obesity: you cannot outrun a bad diet. *Br J Sports Med.* 2015;49:967-8.

Mann KD, Howe LD, Basterfield L ym. Longitudinal study of the associations between change in sedentary behavior and change in adiposity during childhood and adolescence: Gateshead Millennium Study. *Int J Obes.* 2017;41:1042-1047.

Martin CK, Bhapkar M, Pittas AG ym. Effect of calorie restriction on mood, quality of life, sleep, and sexual function in healthy nonobese adults. The CALERIE 2 Randomized Clinical Trial. *JAMA Intern Med.* 2016;176:743-752.

Martins C, Kulseng B, Rehfeld JF, King NA, Blundell JE. Effect of chronic exercise on appetite control in overweight and obese individuals. *Med Sci Sports Exerc.* 2013;45:805-812.

McGuire KA, Ross R. Incidental physical activity and sedentary behavior are not associated with abdominal adipose tissue in inactive adults. *Obesity.* 2012;20:576-82.

McNeil J, Brenner DR, Courneya KS, Friedenreich CM. Dose-response effects of aerobic exercise on energy compensation in postmenopausal women: combined results from two randomized controlled trials. *Int J Obes.* 2017;41:1196-1202.

Mekary RA, Feskanich D, Hu FB, Willett WC, Field AE. Physical activity in relation to long-term maintenance after intentional weight loss in premenopausal women. *Obesity.* 2010;18:167-74.

Melby CL, Paris HL, Foright RM, Peth J. Attenuating the biologic drive for weight regain following weight loss: must what goes down always go back up? *Nutrients.* 2017;9. pii: E468.

Merlotti C, Ceriani V, Morabito A, Pontiroli AE. Subcutaneous fat loss is greater than visceral fat loss with diet and exercise, weight-loss promoting drugs and bariatric surgery: a critical review and meta-analysis. *Int J Obes.* 2017;41:672-682.

Mitchell JA, Bottai M, Park Y, Marshall SJ, Moore SC, Matthews CE. A prospective study of sedentary behavior and changes in the body mass index distribution. *Med Sci Sports Exerc.* 2014;46:2244-52.

Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *New Engl J Med*. 2011;364:2392-404.

Murphy JC, McDaniel JL, Mora K, Villareal DT, Fontana L, Weiss EP. Preferential reductions in intermuscular and visceral adipose tissue with exercise-induced weight loss compared with calorie influence of genetic effects on BMI and waist circumference: a study in young adult twins. *Int J Obes*. 2009 ;33:29-36.

Mustajoki P. Ruokaympäristön muutos selittää pääosan väestöjen lihomisesta. *Duodecim*. 2015;131:1345-52.

Naukkarinen J. Onko tervettä lihavuutta olemassa? *Suomen Lääkäril*. 2017;72:1057.

O'Brien PD, Hinder LM, Callaghan BC, Feldman EL Neurological consequences of obesity. *Lancet Neurol*. 2017;16:465-477.

Ochner CN, Tsai AG, Kushner RF, Wadden TA. Treating obesity seriously: when recommendations for lifestyle change confront biological adaptations. *Lancet Diabetes Endocrinol*. 2015;3:232-4.

Ohkuma T, Hirakawa Y, Nakamura U, Kiyohara Y, Kitazono T, Ninomiya T. Association between eating rate and obesity: a systematic review and meta-analysis. *Int J Obes*. 2015;39:1589-96.

Painter SL, Ahmed R, Hill JO ym. What matters in weight loss? An in-depth analysis of self-monitoring. *J Med Internet Res*. 2017;19:e160.

Parent MC, Alquist JL. Born Fat: The Relations between weight changeability beliefs and health behaviors and physical health. *Health Educ Behav*. 2015 Sep 8. pii: 1090198115602266.

Pate RR, Taverno Ross SE, Liese AD, Dowda M. Associations among physical activity, diet quality, and weight status in US adults. *Med Sci Sports Exerc*. 2015;47:743-50.

Peirson L, Douketis J, Ciliska D, Fitzpatrick-Lewis D, Ali MU, Raina P. Prevention of overweight and obesity in adult populations: a systematic review. *CMAJ Open*. 2014;2:E268-72.

Pihlajamäki J. Geenit ovat osa painonsäätelyä. *Suomen Lääkäril*. 2015;70;105.

Piirtola M, Kaprio J, Waller K ym. Leisure-time physical inactivity and association with body mass index: a Finnish Twin Study with a 35-year follow-up. *Int J Epidemiol*. 2017;46:116-127.

Rillamas-Sun E, LaCroix AZ, Waring ME. Obesity and late-age survival without major disease or disability in older women. *JAMA Intern Med*. 2014;174:98-106.

Robertson C, Archibald D, Avenell A ym. Systematic reviews of and integrated report on the quantitative, qualitative and economic evidence base for the management of obesity in men. *Health Technol Assess*. 2014;18:v-vi, xxiii-xxix, 1-424.

Robinson E, Haynes A, Sutin AR, Daly M. Telling people they are overweight: helpful, harmful or beside the point? *Int J Obes*. 2017;41:1160-1161.

Royal Society for Public Health. Introducing “activity equivalent” calorie labelling to tackle obesity.

http://www.rsph.org.uk/filemanager/root/site_assets/our_work/areas_of_work/activity_calorie_labelling/activity_equivalent_calorie_labelling_paper_rsph.pdf. Viitattu 15.5.2017.

Sahakyan KR, Somers VK, Rodriguez-Escudero JP ym. Normal-weight central obesity: implications for total and cardiovascular mortality. *Ann Intern Med*. 2015;163:827-35.

Sanal E, Ardic F, Kirac S. Effects of aerobic or combined aerobic resistance exercise on body composition in overweight and obese adults:

gender differences. A randomized intervention study. *Eur J Phys Rehabil Med.* 2013;49:1-11.

Shibata AI, Oka K, Sugiyama T, Salmon JO, Dunstan DW, Owen N. Physical activity, television viewing time, and 12-year changes in waist circumference. *Med Sci Sports Exerc.* 2016;48:633-40.

Shiroma EJ, Sesso HD, Lee IM. Physical activity and weight gain prevention in older men. *Int J Obes.* 2012;36:1165-9.

Silventoinen K, Kaprio J. Liikunta auttaa vähentämään geneettisen alttiuden vaikutusta lihavuuteen. *Duodecim.* 2010;126:1031-6.

Sims ST, Larson JC, Lamonte MJ ym. Physical activity and body mass: changes in younger versus older postmenopausal women. *Med Sci Sports Exerc.* 2012;44:89-97.

Stehr MD, von Lengerke T. Preventing weight gain through exercise and physical activity in the elderly: a systematic review. *Maturitas.* 2012;72:13-22.

Stenholm S, Head J, Aalto V ym. Body mass index as a predictor of healthy and disease-free life expectancy between ages 50 and 75: a multicohort study. *Int J Obes.* 2017;41:769-775.

Strandberg TE, Sirola J, Pitkälä KH, Tilvis RS, Strandberg AY, Stenholm S. Association of midlife obesity and cardiovascular risk with old age fragility: a 26-yr follow-up of initially healthy men. *Int J Obes.* 2012;36:1153-57.

The Global Burden of Metabolic Risk Factors for Chronic Diseases Collaboration. Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet.* 2014;383:970-83.

The Lancet Diabetes & Endocrinology. Should we officially recognise obesity as a disease? *Lancet Diabetes & Endocrinology* 2017;5:483

Thomas DM, Martin CK, Lettieri S ym. Can a weight loss of one pound a week be achieved with a 3500-kcal deficit? Commentary on a commonly accepted rule. *Int J Obes.* 2013;37:1611-3.

Thomas DM, Kyle TK, Stanford FC. The gap between expectations and reality of exercise-induced weight loss is associated with discouragement. *Prev Med.* 2015;81:357-60.

Tobias DK, Chen M, Manson JE, Ludwig DS, Willett W, Hu FB. Effect of low-fat diet interventions versus other diet interventions on long-term weight change in adults: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol.* 2015;3:968-79.

Trepanowski JF, Kroeger CM, Barnosky A ym. Effect of alternate-day fasting on weight loss, weight maintenance, and cardioprotection among metabolically healthy obese adults: a randomized clinical trial. *JAMA Intern Med.* 2017;177:930-938.

Zheng Y, Klem ML, Sereika SM, Danford CA, Ewing LJ, Burke LE. Self-weighing in weight management: a systematic literature review. *Obesity.* 2015;23:256-65.

Zheng Y, Manson JE, Yuan C ym. Associations of weight gain from early to middle adulthood with major health outcomes later in life. *JAMA.* 2017;318:255-269.

Veronese N, Facchini S, Stubbs B ym. Weight loss is associated with improvements in cognitive function among overweight and obese people: A systematic review and meta-analysis. *Neurosci Biobehav Rev.* 2017;72:87-94.

Villareal DT, Aguirre L, Gurney AB ym. Aerobic or resistance exercise, or both, in dieting obese older adults. *N Engl J Med.* 2017;376:1943-1955.

Vissers D, Hens W, Taeymans J, Baeyens JP, Poortmans J, Van Gaal L. The effect of exercise on visceral adipose tissue in overweight adults: a systematic review and meta-analysis. *PLoS One*. 2013;8:e56415.

Virtanen KA, Nuutila P. Ihmisen ruskea rasvakudos. *Duodecim*. 2015;131:2075-82.

Wang T, Huang T, Heianza Y ym. Genetic susceptibility, change in physical activity, and long-term weight gain. *Diabetes*. 2017;66:2704-2712.

Westerterp KR. Control of energy expenditure in humans. *Eur J Clin Nutr*. 2017;71:340-344.

Wing RR, Tate DF, Espeland MA ym. The study of novel approaches to weight gain prevention (SNAP) randomized clinical trial. *JAMA Intern Med*. 2016;176:755-762.

Lonkkamurtumat (ks. myös Kaatumiset ja tasapaino ja Osteoporoosi)

Armstrong ME, Cairns BJ, Banks E ym. Different effects of age, adiposity and physical activity on the risk of ankle, wrist and hip fractures in postmenopausal women. *Bone*. 2012;50:1394-400.

Auais MA, Eilayyan O, Mayo NE. Extended exercise rehabilitation after hip fracture improves patients' physical function: a systematic review and meta-analysis. *Phys Ther*. 2012;92:1437-51.

Crandall CJ, Yildiz VO, Wactawski-Wende J ym. Postmenopausal weight change and incidence of fracture: post hoc findings from Women's Health Initiative Observational Study and Clinical Trials. *BMJ*. 2015;350:h25.

Diong J, Allen N, Sherrington C. Structured exercise improves mobility after hip fracture: a meta-analysis with meta-regression. *Br J Sports Med*. 2016;50:346-55.

Donohue K, Hoevenaars R, McEachern J, Zeman E, Mehta S. Home-based multidisciplinary rehabilitation following hip fractures Surgery: what is the evidence? *Rehabil Res Pract*. 2013;2013: Article ID 875968.

Edgren J, Salpakoski A, Sihvonen SE ym. Effects of a home-based physical rehabilitation program on physical disability after hip fracture: a randomized controlled trial. *J Am Med Dir Assoc*. 2015;16:350.e1-7.

El-Khoury F, Cassou B, Charles M-A, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2013;347:f6234.

Englund U, Nordström P, Nilsson J. Physical activity in middle-aged women and hip fracture risk: the UFO study. *Osteoporos Int*. 2011;22:499-505.

Feehan LM, Beck CA, Harris SR, MacIntyre DL, Li LC. Exercise prescription after fragility fracture in older adults: a scoping review. *Osteoporos Int*. 2011;22:1289-322.

Fiatarone Singh MA. Exercise, nutrition and managing hip fracture in older persons. *Curr Opin Clin Nutr Metab Care*. 2014;17:12-24.

Giangregorio LM, MacIntyre NJ, Thabane L, Skidmore CJ, Papaioannou A. Exercise for improving outcomes after osteoporotic vertebral fracture. *Cochrane Database of Systematic Reviews* 2013;1:CD008618.

Haring B, Crandall CJ, Chunyan W ym. Dietary patterns and fractures in postmenopausal women. Results from the Women's Health Initiative. *JAMA Intern Med*. 2016 ;176:645-52.

Kauppi M. Luun lujutta määrittävät ja lonkkamurtumia ennustavat tekijät suomalaisilla aikuisilla. Väitöskirja.

https://www.doria.fi/bitstream/handle/10024/103631/AnnalesD1160Kauppi_Maarit.pdf?sequence=2. Viitattu 23.9.2017.

Kemmler W, Haberle L, von Stengel S. Effects of exercise on fracture reduction in older adults: a systematic review and meta-analysis. *Osteoporosis Int.* 2013;24:1937-1950.

Latham NK, Harris BA, Bean JF ym. Effect of a home-based exercise program on functional recovery following rehabilitation after hip fracture: a randomized clinical trial. *JAMA* 2014;311:700-8.

Lonkkamurtuma - Käypä hoito. 16.10.2017.

<http://www.kaypahoito.fi/web/kh/suositukset/suositus?id=hoi50040>.

Viitattu 14.11.2017.

Morseth B, Ahmed LA, Bjørnerem Å. Leisure time physical activity and risk of non-vertebral fracture in men and women aged 55 years and older: the Tromsø Study. *Eur J Epidemiol.* 2012;27:463-71.

Määttä M, Terho E, Jokinen H ym. Lifestyle factors and site-specific risk of hip fracture in community dwelling older women--a 13-year prospective population-based cohort study. *BMC Musculoskelet Disord.* 2012;13:173.

Nikander R, Gagnon C, Dunstan DW ym, Frequent walking, but not total physical activity, is associated with increased fracture incidence: a 5-year follow-up of an Australian population-based prospective study (AusDiab). *J Bone Miner Res.* 2011;26:1638-47

Rikkonen T. Physical activity and muscle strength as predictors of bone loss and fractures in perimenopausal women. *Dissertations in Health Sciences* 146, 2012. Itä-Suomen yliopisto.

Salpakoski A, Törmäkangas T, Edgren J ym. Effects of a multicomponent home-based physical rehabilitation program on mobility recovery after hip fracture: a randomized controlled trial. *J Am Med Dir Assoc.* 2014 ;15:361-8.

Sherrington C, Tiedemann A, Cameron ID. Physical exercise after hip fracture: an evidence overview. *Eur J Phys Rehabil Med.* 2011;47:297-307.

Stott D, Handoll H. Rehabilitation of older people after hip (proximal femur) fracture. *Cochrane Database Syst Rev.* 2011;8:CD00023.

Stattin K, Michaëlsson K, Larsson SC, Wolk A, Byberg L. Leisure-time physical activity and risk of fracture: a cohort study of 66,940 men and women. *J Bone Miner Res.* 2017;32:1599-1606.

Turunen K, Salpakoski A, Edgren J ym. Physical activity after a hip fracture: effect of a multicomponent home-based rehabilitation program - a secondary analysis of a randomized controlled trial. *Arch Phys Med Rehabil.* 2017 Jan 27. pii: S0003-9993(17)30034-5.

Zhao R, Feng F, Wang X. Exercise interventions and prevention of fall-related fractures in older people: a meta-analysis of randomized controlled trials. *Int J Epidemiol.* 2017;46:149-161.

Niskakipu

Alpayci M, İltis S. Isometric exercise for the cervical extensors can help restore physiological lordosis and reduce neck pain: a randomized controlled trial. *Am J Phys Med Rehabil.* 2017;96:621-626.

Andersen CH, Andersen LL, Gram B ym. Influence of frequency and duration of strength training for effective management of neck and shoulder pain: a randomised controlled trial. *Br J Sports Med.* 2012 ;46:1004-10.

Andersen LL, Saervoll CA, Mortensen OS, Poulsen OM, Hannerz H, Zebis MK. Effectiveness of small daily amounts of progressive resistance training for frequent neck/shoulder pain: randomized controlled trial. *Pain.* 2011;152:440-46.

Arokoski, J, Laimi K. Nuoren niskakipu on yleinen vaiva. *Suomen Lääkäril.* 2014;69:879-884.

Arokoski J, Karppinen J, Kankaanpää M, Kaukinen P, Lami K. Aikuisen kipeä niska. *Duodecim*. 2014;130:2099-107.

Bertozzi L, Gardenghi I, Turoni F ym. Effect of therapeutic exercise on pain and disability in the management of chronic nonspecific neck pain: systematic review and meta-analysis of randomized trials. *Phys Ther*. 2013;93:1026-36.

Bervoets DC, Luijsterburg PA, Alessie JJ, Buijs MJ, Verhagen AP. Massage therapy has short-term benefits for people with common musculoskeletal disorders compared to no treatment: a systematic review. *J Physiother*. 2015 ;61:106-16.

Bishop MD, Mintken PE, Bialosky JE, Cleland JA. Patient expectations of benefit from interventions for neck pain and resulting influence on outcomes. *J Orthop Sports Phys Ther*. 2013;43:457-65.

Bronfort G, Evans R, Anderson AV, Svendsen KH, Bracha Y, Grimm RH. Spinal manipulation, medication, or home exercise with advice for acute and subacute neck pain: a randomized trial. *Ann Intern Med*. 2012;156:1-10.

Bryans R, Decina P, Descarreaux M. Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manipulative Physiol Ther*. 2014;37:42-63.

Carroll LJ, Hogg-Johnson S, Côté P ym. Course and prognostic factors for neck pain in workers: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*. 2008;33:S93-100.

Cheng YH, Huang GC. Efficacy of massage therapy on pain and dysfunction in patients with neck pain: a systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2014;2014:204360.

Côté P, Wong JJ, Sutton D ym. Management of neck pain and associated disorders: A clinical practice guideline from the Ontario Protocol for

Traffic Injury Management (OPTIMa) Collaboration. *Eur Spine J.* 2016;25:2000-22.

Crawford C, Boyd C, Paat CF ym. The impact of massage therapy on function in pain populations – a systematic review and meta-analysis of randomized controlled trials: Part I, Patients experiencing pain in the general population. *Pain Med.* 2016 May 10. pii: pnw099.

Furlan AD, Imamura M, Dryden T, Irvin E. Massage for low-back pain. *Cochrane Database Syst Rev.* 2015;9:CD001929.

Furlan AD Yazdi F, Tsertsvadze A ym. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med.* 2012;2012:953139.

Garra G, Singer AJ, Leno R ym. Heat or cold packs for neck and back strain: a randomized controlled trial of efficacy. *Acad Emerg Med.* 2010;17:484-9.

Gram B, Andersen C, Zebis MK ym. Effect of training supervision on effectiveness of strength training for reducing neck/shoulder pain and headache in office workers: cluster randomized controlled trial. *Biomed Res Int.* 2014;2014:693013.

Gross A, Kay TM, Paquin JP ym. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev.* 2015 ;1:CD004250.

Gross A, Miller J, D'Sylva J ym. Manipulation or mobilisation for neck pain. *Cochrane Database Syst Rev.* 2015;9:CD004249

Hallman DM, Birk Jørgensen M, Holtermann A. Objectively measured physical activity and 12-month trajectories of neck-shoulder pain in workers: A prospective study in DPHACTO. *Scand J Public Health.* 2017;45:288-298.

Hagen KB, Dagfinrud H, Moe RH ym. Exercise therapy for bone and muscle health: an overview of systematic reviews. *BMC Medicine* 2012;10:167.

Hoe VC, Urquhart DM, Kelsall HL, Sim MR. Ergonomic design and training for preventing work-related musculoskeletal disorders of the upper limb and neck in adults. *Cochrane Database Syst Rev.* 2012;8:CD008570.

Jay K, Schraefel M, Andersen CH, Ebbesen FS ym. Effect of brief daily resistance training on rapid force development in painful neck and shoulder muscles: randomized controlled trial. *Clin Physiol Funct Imaging.* 2013;33:386-92.

Karlsson L, Gerdle B, Takala EP, Andersson G, Larsson B. Associations between psychological factors and the effect of home-based physical exercise in women with chronic neck and shoulder pain. *SAGE Open Med.* 2016;4:2050312116668933.

Kauranen K. Epäspesifinen niskakipu, s. 69-77. Teoksessa Kauranen K. *Fysioterapeutin käsikirja*, Sanoma Pro 2017, Helsinki..

Kjaer P, Kongsted A, Hartvigsen J ym. National clinical guidelines for non-surgical treatment of patients with recent onset neck pain or cervical radiculopathy. *Eur Spine J.* 2017 May 18.

Kong LJ, Zhan HS, Cheng YW, Yuan WA, Chen B, Fang M. Massage therapy for neck and shoulder pain: a systematic review and meta-analysis. *Evid Based Complement Alternat Med.* 2013;2013:613279.

Lee KW, Kim WH. Effect of thoracic manipulation and deep craniocervical flexor training on pain, mobility, strength, and disability of the neck of patients with chronic nonspecific neck pain: a randomized clinical trial. *J Phys Ther Sci.* 2016;28:175-80.

Li X, Lin C, Liu C ym. Comparison of the effectiveness of resistance training in women with chronic computer-related neck pain: a randomized controlled study. *Int Arch Occup Environ Health.* 2017 May 20

Lin LY, Wang RH. Effectiveness of a neck stretching intervention on nurses' primary headaches. *Workplace Health Saf.* 2015;63:100-6.

Lowe BD, Dick RB. Workplace exercise for control of occupational neck/shoulder disorders: a review of prospective studies. *Environ Health Insights.* 2015;8(Suppl 1):75-95.

Mortensen P, Larsen AI, Zebis MK ym. Lasting effects of workplace strength training for neck/ shoulder/arm pain among laboratory technicians: natural experiment with 3-year follow-up. *Biomed Res Int.* 2014;2014:845851.

Nilsen TI, Holtermann A, Mork PJ. Physical exercise, body mass index, and risk of chronic pain in the low back and neck/shoulders: longitudinal data from the Nord-Trøndelag Health Study. *Am J Epidemiol* 2011;174:267-73 .

Niskakipu (aikuiset). Käypä hoito-suositus. 16.01.2017. Suomalaisen Lääkäriseuran Duodecimin, Societas Medicinae Physicalis et Rehabilitationis Fenniae ry:n ja Suomen Yleislääketieteen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi20010>.

Viitattu 15.11.2017.

O'Riordan C, Clifford A, Van De Ven P, Nelson J. Chronic neck pain and exercise interventions: frequency, intensity, time, and type principle. *Arch Phys Med Rehabil.* 2014;95:770-83.

Paanalahti K, Holm LW, Nordin M, Asker M, Lyander J, Skillgate E. Adverse events after manual therapy among patients seeking care for neck and/or back pain: a randomized controlled trial. *BMC Musculoskelet Disord.* 2014;15:77.

Paanalahti K, Holm LW, Nordin M ym. Three combinations of manual therapy techniques within naprapathy in the treatment of neck and/or back

pain: a randomized controlled trial. *BMC Musculoskelet Disord.* 2016;17:176.

Paksaichol A, Janwantanakul P, Purepong N, Pensri P, van der Beek AJ. Office workers' risk factors for the development of non-specific neck pain: a systematic review of prospective cohort studies. *Occup Environ Med.* 2012;69:610-8.

Patel KC, Gross A, Graham N ym. Massage for mechanical neck disorders. *Cochrane Database Syst Rev.* 2012;9:CD004871.

Pedersen MT, Andersen CH, Zebis MK, Sjøgaard G, Andersen LL. Implementation of specific strength training among industrial laboratory technicians: long-term effects on back, neck and upper extremity pain. *BMC Musculoskelet Disord.* 2013;14:287.

Poppendieck W, Wegmann M, Ferrauti A, Kellmann M, Pfeiffer M, Meyer T. Massage and performance recovery: a meta-analytical review. *Sports Med.* 2016;46:183-204.

Salo P, Ylönen-Käyrä N, Häkkinen A, Kautiainen H, Mälkiä E, Ylinen J. Effects of long-term home-based exercise on health-related quality of life in patients with chronic neck pain: a randomized study with a 1-year follow-up. *Disabil Rehabil.* 2012;34:1971-77.

SBU. Preventiva insatser vid akut smärta från rygg och nacke – Effekter av fysisk träning, manuell behandling och beteendepåverkande åtgärder. Stockholm: Statens beredning för medicinsk och social utvärdering (SBU). 2016.<http://www.sbu.se/sv/publikationer/SBU-utvarderar/preventiva-insatser-vid-akut-smarta-fran-rygg-och-nacke--effekter-av-fysisk-traning-manuell-behandling-och-beteendepaverkande-atgarder/3>. Viitattu 21.10.2017.

Southerst D, Nordin MC, Côté P ym. Is exercise effective for the management of neck pain and associated disorders or whiplash-associated

disorders? A systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Spine J.* 2016;16:1503-23.

Suni JH, Rinne M, Tokola K, Mänttari A, Vasankari T. Effectiveness of a standardised exercise programme for recurrent neck and low back pain: a multicentre, randomised, two-arm, parallel group trial across 34 fitness clubs in Finland. *BMJ Open Sport Exerc Med.* 2017;3:e000233.

Trinh KV, Graham N, Gross AR, Goldsmith CH, Wang E, Cameron ID, Kay T; Cervical Overview Group.. Acupuncture for neck disorders. *Cochrane Database Syst Rev.* 2016;5:CD004870.

Zaproudina N, Hänninen OO, Airaksinen O. Effectiveness of traditional bone setting in chronic neck pain: randomized clinical trial. *J Manipulative Physiol Ther.* 2007;30:432-7.

Zebis MK, Andersen LL, Pedersen MT ym. Implementation of neck/shoulder exercises for pain relief among industrial workers: a randomized controlled trial. *BMC Musculoskelet Disord.* 2011;12:205.

Varatharajan S, Ferguson B, Chrobak K ym. Are non-invasive interventions effective for the management of headaches associated with neck pain? An update of the Bone and Joint Decade Task Force on Neck Pain and Its Associated Disorders by the Ontario Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Eur Spine J.* 2016;25:1971-99.

Wong JJ, Shearer HM, Mior S ym. Are manual therapies, passive physical modalities, or acupuncture effective for the management of patients with whiplash-associated disorders or neck pain and associated disorders? An update of the Bone and Joint Decade Task Force on Neck Pain and Its Associated Disorders by the OPTIMa collaboration. *Spine J.* 2016;16:1598-1630.

Yuan QL, Guo TM, Liu L, Sun F, Zhang YG. Traditional Chinese medicine for neck pain and low back pain: a systematic review and meta-analysis. *PLoS One*. 2015;10:e0117146.

Ylinen J, Nikander R. Harjoittelun vaikuttavuus ja toteutus kroonisen niskakipupotilaankuntoutuksessa. *Suomen Lääkäril*. 2014;69:2475-61.

Nivelreuma

Agca R, Heslinga SC, Van Halm VP, Nurmohamed MT. Atherosclerotic cardiovascular disease in patients with chronic inflammatory joint disorders. *Heart*. 2016;102:790-5.

Al-Qubaeissy KY, Fatoye FA, Goodwin PC, Yohannes AM. The effectiveness of hydrotherapy in the management of rheumatoid arthritis: a systematic review. *Musculoskeletal Care*. 2013;11:3-18.

Baillet A, Zeboulon N, Gossec L ym. Efficacy of cardiorespiratory aerobic exercise in rheumatoid arthritis: meta-analysis of randomized controlled trials. *Arthritis Care Res*. 2010;62:984-92.

Baillet A, Vaillant M, Guinot M, Juvin R, Gaudin P. Efficacy of resistance exercises in rheumatoid arthritis: meta-analysis of randomized controlled trials. *Rheumatology*. 2012;51:519-27.

Burr J, Shephard R, Cornish S, Vatanparast H, Chilibeck P. Arthritis, osteoporosis, and low back pain: evidence-based clinical risk assessment for physical activity and exercise clearance. *Can Fam Physician*. 2012;58:59-62.

Cameron M, Gagnier JJ, Chrubasik S. Herbal therapy for treating rheumatoid arthritis. *Cochrane Database Syst Rev*. 2011;2:CD002948.

Casimiro L, Barnsley L, Brosseau L ym. Acupuncture and electroacupuncture for the treatment of rheumatoid arthritis. *Cochrane Database Syst Rev*. 2005;4:CD003788.

Chang K, Yang SM, Kim SH, Han KH, Park SJ, Shin JI. Smoking and rheumatoid arthritis. *Int J Mol Sci.* 2014 3;15:22279-95.

Cooney JK, Law R-J, Matscke V ym. Benefits of exercise in rheumatoid arthritis. *J Aging Res.* 2011, Article ID 681640.

Cramer H, Lauche R, Langhorst J, Dobos G. Yoga for rheumatic diseases: a systematic review. *Rheumatology.* 2013;52:2025-30.

Cramp F, Hewlett S, Almeida C ym. Non-pharmacological interventions for fatigue in rheumatoid arthritis. *Cochrane Database Syst Rev.* 2013;8:CD008322.

Di Giuseppe D, Bottai M, Askling J, Wolk A. Physical activity and risk of rheumatoid arthritis in women: a population-based prospective study. *Arthritis Res Ther.* 2015;17:40.

Di Giuseppe D, Crippa A, Orsini N, Wolk A. Fish consumption and risk of rheumatoid arthritis: a dose-response meta-analysis. *Arthritis Res Ther.* 2014;16:446.

Di Giuseppe D, Discacciati A, Orsini N, Wolk A. Cigarette smoking and risk of rheumatoid arthritis: a dose-response meta-analysis. *Arthritis Res Ther.* 2014;16:R61.

Di Giuseppe D, Wallin A, Bottai M, Askling J, Wolk A. Long-term intake of dietary long-chain n-3 polyunsaturated fatty acids and risk of rheumatoid arthritis: a prospective cohort study of women. *Ann Rheum Dis.* 2014;73:1949-53.

Feldthusen C, Dean E, Forsblad-d'Elia H, Mannerkorpi K. Effects of person-centered physical therapy on fatigue-related variables in persons with rheumatoid arthritis: a randomized controlled trial. *Arch Phys Med Rehabil.* 2016;97:26-36.

Feldthusen C, Grimby-Ekman A, Forsblad-d'Elia H, Jacobsson L, Mannerkorpi K. Explanatory factors and predictors of fatigue in persons

with rheumatoid arthritis: A longitudinal study. *J Rehabil Med*. 2016;48:469-76.

Fenton SA, Veldhuijzen van Zanten JJ, Duda JL, Metsios GS, Kitas GD. Sedentary behaviour in rheumatoid arthritis: definition, measurement and implications for health. *Rheumatology*. 2017 Apr 7.

Fenton SAM, Veldhuijzen van Zanten JJCS, Kitas GD, Duda JL, Rouse PC, Yu CA, Metsios GS. Sedentary behaviour is associated with increased long-term cardiovascular risk in patients with rheumatoid arthritis independently of moderate-to-vigorous physical activity. *BMC Musculoskelet Disord*. 2017;18:131.

Guillot X, Tordi N, Mourot L ym. Cryotherapy in inflammatory rheumatic diseases: a systematic review. *Expert Rev Clin Immunol*. 2014;10:281-94.

Hagen KB, Byfuglien MG, Falzon L, Olsen SU, Smedslund G. Dietary interventions for rheumatoid arthritis. *Cochrane Database Syst Rev*. 2009;1:CD006400.

Hagen KB, Dagfinrud H, Moe RH ym. Exercise therapy for bone and muscle health: an overview of systematic reviews. *BMC Medicine*. 2012;10:167.

Hurkmans E, van der Giesen FJ, Vliet Vlieland TP,ym. Dynamic exercise programs (aerobic capacity and/or muscle strength training) in patients with rheumatoid arthritis. *Cochrane Database Syst Rev*. 2009;4:CD006853.

Jin Z, Xiang C, Cai Q, Wei X, He J. Alcohol consumption as a preventive factor for developing rheumatoid arthritis: a dose-response meta-analysis of prospective studies. *Ann Rheum Dis*. 2014;73:1962-7.

Julkunen H, Eklund K. Nivelituloehdusten diagnostiikka. *Duodecim* 2017;133:1293-301.

Karlson EW, Shacick NA, Cook NR, Buring JE, Lee IM. Vitamin E in the primary prevention of rheumatoid arthritis: the Women's Health Study. *Arthritis Rheu.* 2008;59:1589-95.

Kelley GA, Kelley KS, Hootman JM, Jones DL. Effects of community-deliverable exercise on pain and physical function in adults with arthritis and other rheumatic diseases: a meta-analysis. *Arthritis Care Res.* 2011;63:79-93.

Kerola A. Epidemiology of comorbidities in early rheumatoid arthritis: with emphasis on cardiovascular disease. Väitöskirja.
<http://urn.fi/URN:ISBN:978-951-51-0900-2>. Viitattu 12.3.2017.

Koivuniemi R. Leirisalo-Repo M. Tulehduksellisiin reumasairauksiin liittyy suurentunut sydän- ja verisuonitautien riski. *Suomen Lääkäril.* 2015;70:711-15.

Lahiri M, Morgan C, Symmons DP, Bruce IN. Modifiable risk factors for RA: prevention, better than cure? *Rheumatology.* 2012;51:499-512.

Lahiri M, Luben RN, Morgan C ym. Using lifestyle factors to identify individuals at higher risk of inflammatory polyarthritis (results from the European Prospective Investigation of Cancer-Norfolk and the Norfolk Arthritis Register--the EPIC-2-NOAR Study). *Ann Rheum Dis.* 2014;73:219-26.

Lamb SE, Williamson EM, Heine PJ ym. Exercises to improve function of the rheumatoid hand (SARAH): a randomised controlled trial. *Lancet.* 2015;385(9966):421-9.

Lemmey AB, Wikkiams SL, Marcora SM, Jones L, Maddison PJ. Are the benefits of a high-intensity progressive resistance training program sustained in rheumatoid arthritis patients? A 3-year follow-up study. *Arthritis Care Res.* 2012;64:71-5.

Lu, B, Hiraki LT, Sparks JA ym. Being overweight or obese and risk of developing rheumatoid arthritis among women: a prospective cohort study. *Ann Rheum Dis*. 2014 Jul 23. pii: annrheumdis-2014-205459.

Moonaz SH, Bingham CO 3rd, Wissow L, Bartlett SJ. Yoga in sedentary adults with arthritis: effects of a randomized controlled pragmatic trial. *J Rheumatol*. 2015;42:1194-202.

Nivelreuma. Käypä hoito -suositus. 08.06.2015. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Reumatologisen yhdistyksen asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi21010>

Viitattu 12.10.2017.

Qin B, Yang M, Fu H ym. Body mass index and the risk of rheumatoid arthritis: a systematic review and dose-response meta-analysis. *Arthritis Res Ther*. 2015;17:86

Robinson V, Brosseau L, Casimiro L. Thermotherapy for treating rheumatoid arthritis. *Cochrane Database Syst Rev*. 2002;1:CD002826.

Rongen-van Dartel SA, Repping-Wuts H, Flendrie M ym. Effect of aerobic exercise training on fatigue in rheumatoid arthritis: a meta-analysis. *Arthritis Care Res*. 2015;67:1054-62.

Scott IC, Tan R, Stahl D, Steer S, Lewis CM, Cope AP. The protective effect of alcohol on developing rheumatoid arthritis: a systematic review and meta-analysis. *Rheumatology*. 2013;52:856-67.

Sharif K, Watad A, Bragazzi N, Lichtbroun M, Amital H, Shoenfeld Y. Physical activity and autoimmune diseases: Get moving and manage the disease. *Autoimmun Rev*. 2018;17:53-72.

Smedslund G1, Byfuglien MG, Olsen SU, Hagen KB. Effectiveness and safety of dietary interventions for rheumatoid arthritis: a systematic review of randomized controlled trials. *J Am Diet Assoc*. 2010;110:727-35.

Smith BE, Hendrick P, Smith TO. Should exercises be painful in the management of chronic musculoskeletal pain? A systematic review and meta-analysis. *Br J Sports Med.* 2017;51:1679-1687.

Strasser B, Leeb G, Strehlow C, Schobersberger W, Haber P, Cauza E. The effects of strength and endurance training in patients with rheumatoid arthritis. *Clin Rheumatol.* 2011;30:623-32.

Veldhuijzen van Zanten JJ, Rouse PC, Hale ED, Ntoumanis N, Metsios GS, Duda JL, Kitas GD. Perceived barriers, facilitators and benefits for regular physical activity and exercise in patients with rheumatoid arthritis: a review of the literature. *Sports Med.* 2015 ;45:1401-12.

Verhagen AP, Bierma-Zeinstra SM, Boers M, Cardoso JR, Lambeck J, de Bie R, de Vet HC. Balneotherapy (or spa therapy) for rheumatoid arthritis. *Cochrane Database Syst Rev.* 2015;4:CD000518.

Vliet Vlieland TP, van den Ende CH. Nonpharmacological treatment of rheumatoid arthritis. *Curr Opin Rheumatol.* 2011;23:259-64.

Nivelriikko

Abbott JH, Robertson MC, Chapple C ym. Manual therapy, exercise therapy, or both, in addition to usual care, for osteoarthritis of the hip or knee: a randomized controlled trial. 1: clinical effectiveness. *Osteoarthritis Cartilage.* 2013;21:525-534.

Ageberg E, Engström G, Gerhardsson de Verdier, Rollof J, Roos EM, Lohmander LS. Effect of leisure time physical activity on severe knee or hip osteoarthritis leading to total joint replacement: a population-based prospective cohort study. *BMC Musculoskelet Disord.* 2012;13:73.

Allen KD, Golightly YM. Epidemiology of osteoarthritis: state of the evidence. *Curr Opin Rheumatol.* 2015; 27:276–283.

Anwer S, Alghadir A, Brismée JM. Effect of home exercise program in patients with knee osteoarthritis: a systematic review and meta-analysis. *J Geriatr Phys Ther.* 2016;39:38-48.

Apold H, Meyer HE, Nordsletten L, Furnes O, Baste V, Flugsrud GB. Weight gain and the risk of knee replacement due to primary osteoarthritis: a population based, prospective cohort study of 225,908 individuals. *Osteoarthritis Cartilage.* 2014;22:652-8.

Apold H, Meyer HE, Nordsletten L, Furnes O, Baste V, Flugsrud GB. Risk factors for knee replacement due to primary osteoarthritis, a population based, prospective cohort study of 315,495 individuals. *BMC Musculoskelet Disord.* 2014; 15: 217.

Atukorala I, Makovey J, Lawler L, Messier SP, Bennell K, Hunter DJ. Is there a dose response relationship between weight loss and symptom improvement in persons with knee osteoarthritis? *Arthritis Care Res.* 2016;68:1106-14.

Barbour KE, Hootman JM, Helmick CG ym. Meeting physical activity guidelines and the risk of incident knee osteoarthritis: a population-based prospective cohort study. *Arthritis Care Res.* 2014;66:139-46.

Bartels EM, Juhl CB, Christensen R ym. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev.* 2016;3:CD005523.

Bastick AN, Runhaar J, Belo JN, Bierma-Zeinstra SM. Prognostic factors for progression of clinical osteoarthritis of the knee: a systematic review of observational studies. *Arthritis Res Ther.* 2015;17:152.

Bastick AN, Verkleij SP, Damen J, Wesseling J, Hilberdink WK, Bindels PJ, Bierma-Zeinstra SM. Defining hip pain trajectories in early symptomatic hip osteoarthritis – 5 year results from a nationwide prospective cohort study (CHECK). *Osteoarthritis Cartilage.* 2016;24:768-75.

Bastick AN, Wesseling J, Damen J ym. Defining knee pain trajectories in early symptomatic knee osteoarthritis in primary care: 5-year results from a nationwide prospective cohort study (CHECK). *Br J Gen Pract.* 2016;66:e32-9.

Batsis JA, Zbehlik AJ, Barre LK, Bynum JP, Pidgeon D, Bartels SJ. Impact of obesity on disability, function, and physical activity: data from the Osteoarthritis Initiative. *Scand J Rheumatol.* 2015;44:495-502.

Batterham SI, Heywood S, Keating JL. Systematic review and meta-analysis comparing land and aquatic exercise for people with hip or knee arthritis on function, mobility and other health outcomes. *BMC Musculoskelet Disord.* 2011;12:123.

Bennell KL, Egerton T, Martin J ym. Effect of physical therapy on pain and function in patients with hip osteoarthritis. *JAMA.* 2014;311:1987-1997.

Bieler T, Siersma V, Magnusson SP, Kjaer M, Christensen HE, Beyer N. In hip osteoarthritis, Nordic walking is superior to strength training and home-based exercise for improving function. *Scand J Med Sci Sports.* 2017 ;27:873-886.

Brosseau L, Wells GA, Pugh AG. Ottawa Panel evidence-based clinical practice guidelines for therapeutic exercise in the management of hip osteoarthritis. *Clin Rehabil.* 2015 Sep 23. pii: 0269215515606198.

Buttgereit F, Burmester GR, Bijlsma JW. Non-surgical management of knee osteoarthritis: where are we now and where do we need to go? *RMD Open.* 2015;1:e000027.

Burr J , Shephard R, Cornish S, Vatanparast H, Chilibeck P. Arthritis, osteoporosis, and low back pain: evidence-based clinical risk assessment for physical activity and exercise clearance. *Can Fam Physician.* 2012;58:59-62.

Cheung C, Wyman JF, Bronas U, McCarthy T, Rudser K, Mathiason MA. Managing knee osteoarthritis with yoga or aerobic/strengthening exercise programs in older adults: a pilot randomized controlled trial. *Rheumatol Int.* 2017;37:389-398.

Collins JE, Katz JN, Dervan EE, Losina E. Trajectories and risk profiles of pain in persons with radiographic, symptomatic knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthritis Cartilage.* 2014 ;22:622-30.

Coulter C, Perriman DM, Neeman TM, Smith PN, Scarvell JM. Supervised or unsupervised rehabilitation after total hip replacement provides similar improvements for patients: a randomized controlled trial. *Arch Phys Med Rehabil.* 2017;98:2253-2264.

Cotofana S, Ring-Dimitriou S, Hudelmaier M ym. Effects of exercise intervention on knee morphology in middle-aged women: a longitudinal analysis using magnetic resonance imaging. *Cells Tissues Organs.* 2010;192:64-72.

Culvenor AG, Felson DT, Niu J ym. Thigh muscle specific strength and the risk of incident knee osteoarthritis: The influence of sex and greater body mass index. *Arthritis Care Res.* 2017. Feb 8. doi: 10.1002/acr.23182.

de Rooij M, van der Leeden M, Heymans MW. Prognosis of pain and physical functioning in patients with knee osteoarthritis: a systematic review and meta-analysis. *Arthritis Care Res.* 2016 ;68:481-92.

Ding C, Stannus O, Cicuttini F, Antony B, Jones G. Body fat is associated with increased and lean mass with decreased knee cartilage loss in older adults: a prospective cohort study. *Int J Obesity* 2013;37:822-27.

Dunlop DD, Song J, Semanik PA ym. Relation of physical activity time to incident disability in community dwelling adults with or at risk of knee arthritis: prospective cohort study. *BMJ.* 2014;348:g2472.

- Driban JB, Hootman JM, Sitler MR, Harris K, Cattano NM. Is participation in certain sports associated with knee osteoarthritis? A systematic review. *J Athl Train*. 2017;52:497-506.
- Dubé CE, Liu SH, Driban JB, McAlindon TE, Eaton CB, Lapane KL. The relationship between smoking and knee osteoarthritis in the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2016;24:465-72.
- Felson DT, Hodgson R. Identifying and treating preclinical and early osteoarthritis. *Rheum Dis Clin North Am*. 2014;40:699-710.
- Felson DT, Zhang Y. Smoking and osteoarthritis: a review of the evidence and its implications. *Osteoarthritis Cartilage*. 2015;23:331-3.
- Fernandes L, Hagen KB, Bijlsma JW ym. EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis*. 2013;72:1125-35.
- Fernandopulle S, Perry M, Manlapaz D, Jayakaran P. Effect of land-based generic physical activity interventions on pain, physical function, and physical performance in hip and knee osteoarthritis: a Systematic review and meta-analysis. *Am J Phys Med Rehabil*. 2017;96:773-792.
- Fransen M, McConnell S, Hernandez-Molina G, Reichenbach S. Exercise for osteoarthritis of the hip. *Cochrane Database Syst Rev*. 2014;4:CD007912.
- Fransen M, McConnell S, Harmer AR, Van der Esch M, Simic M, Bennell KL. Exercise for osteoarthritis of the knee: a Cochrane systematic review. *Br J Sports Med*. 2015;49:1554-7.
- Gallagher B, Tjoumakaris FP, Harwood MI, Good RP, Cicotti MG, Freedman KB. Chondroprotection and the prevention of osteoarthritis progression of the knee: a systematic review of treatment agents. *Am J Sports Med*. 2015;43:734-44.

Gersing AS, Solka M, Joseph GB ym. Progression of cartilage degeneration and clinical symptoms in obese and overweight individuals is dependent on the amount of weight loss: 48-month data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2016;24:1126-34.

Geusens PP, van den Bergh JP. Osteoporosis and osteoarthritis: shared mechanisms and epidemiology. *Curr Opin Rheumatol*. 2016;28:97-103.

Glyn-Jones S, Palmer AJ, Agricola R, Price AJ, Vincent TL, Weinans H, Carr AJ. Osteoarthritis. *Lancet*. 2015;386(9991):376-87.

Golightly YM, Allen KD, Caine DJ. A comprehensive review of the effectiveness of different exercise programs for patients with osteoarthritis. *Phys Sportsmed*. 2012;40:52-65.

Gouttebauge V, Inklaar H, Backx F, Kerkhoffs G. Prevalence of osteoarthritis in former elite athletes: a systematic overview of the recent literature. *Rheumatol Int*. 2015;35:405-18.

Gudbergesen H, Boesen M, Lohmander LS ym. Weight loss is effective for symptomatic relief in obese subjects with knee osteoarthritis independently of joint damage severity assessed by high-field MRI and radiography. *Osteoarthritis Cartilage*. 2012;20:495-502.

Haaz S, Bartlett SJ. Yoga for arthritis: a scoping review. *Rheum Dis Clin North Am*. 2011;37:33-46.

Henriksen M, Klokke L, Graven-Nielsen T ym. Association of exercise therapy and reduction of pain sensitivity in patients with knee osteoarthritis: a randomized controlled trial. *Arthritis Care Res*. 2014;66:1836-43.

Hochberg MC, Altman RD, April KT ym. American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res*. 2012 ;64:465-74.

Holla JF, van der Leeden M, Heymans MW ym. Three trajectories of activity limitations in early symptomatic knee osteoarthritis: a 5-year follow-up study. *Ann Rheum Dis*. 2014;73:1369-75.

Holla JF, van der Leeden M, Knol DL ym. Predictors and outcome of pain-related avoidance of activities in persons with early symptomatic knee osteoarthritis: A 5-year follow-up study. *Arthritis Care Res*. 2014; 66:802–809.

Hunter DJ. Viscosupplementation for osteoarthritis of the knee. *N Engl J Med*. 2015;372:2570.

Iosifidis MI, Tsarouhas A, Fylaktou A. Lower limb clinical and radiographic osteoarthritis in former elite male athletes. *Knee Surg Sports Traumatol Arthrosc*. 2015;23:2528-35.

Jansen MJ, Viechtbauer W, Lenssen AF, Hendriks EJ, de Bie RA. Strength training alone, exercise therapy alone, and exercise therapy with passive manual mobilisation each reduce pain and disability in people with knee osteoarthritis: a systematic review. *J Physiother*. 2011;57:11-20.

Jones G, Bennell K, Cicuttini FM. Effect of physical activity on cartilage development in healthy kids. *Br J Sports Med*. 2003;37:382-3.

Jorge RT, Souza MC, Chiari A, Jones A, Fernandes Ada R, Lombardi Júnior I, Natour J. Progressive resistance exercise in women with osteoarthritis of the knee: a randomized controlled trial. *Clin Rehabil*. 2015 ;29:234-43.

Juhakoski R. Hip osteoarthritis; risk factors and the effects of exercise therapy (Lonkkanivelriikon riskitekijät ja terapeuttisen harjoittelun vaikutukset lonkkanivelrikkopotilaan toimintakykyyn). Itä-Suomen yliopisto 2013. ISBN (pdf): 978-952-61-1104-9.

Juhakoski R, Arokoski J. Lonkkanivelriikon konservatiivinen hoito. *Suomen Lääkäril*. 2014;69:2700-6.

Juhl C, Christensen R, Roos EM, Zhang W, Lund H. Impact of exercise type and dose on pain and disability in knee osteoarthritis: A systematic review and meta-regression analysis of randomized controlled trials. *Arthr Rheumatol.* 2014;66:622-36.

Kang JW, Lee MS, Posadzki P, Ernst E. T'ai chi for the treatment of osteoarthritis: a systematic review and meta-analysis. *BMJ Open.* 2011;1:e000035.

Karlson EW, Shadick NA, Cook NR, Buring JE, Lee IM. Vitamin E in the primary prevention of rheumatoid arthritis: the Women's Health Study. *Arthritis Rheum.* 2008;59:1589-95.

Kelley GA, Kelley KS, Hootman JM, Jones DL. Effects of community-deliverable exercise on pain and physical function in adults with arthritis and other rheumatic diseases: a meta-analysis. *Arthritis Care Res.* 2011;63:79-93.

Kim C, Nevitt MC, Niu J ym. Association of hip pain with radiographic evidence of hip osteoarthritis: diagnostic test study. *BMJ.* 2015;351:h5983.

Koli J, Multanen J, Kujala UM ym. Effects of exercise on patellar cartilage in women with mild knee osteoarthritis. *Med Sci Sports Exerc.* 2015;47:1767-74.

Konstari S, Kaila-Kangas L, Jääskeläinen T ym. Serum 25-hydroxyvitamin D and the risk of knee and hip osteoarthritis leading to hospitalization: a cohort study of 55274 Finns. *Rheumatology* 2014;53:1778--82.

Kristensen J, Franklyn-Miller A. Resistance training in musculoskeletal rehabilitation: a literature review. *Br J Sports Med.* 2012;46:719-26.

Kuru Çolak T, Kavlak B, Aydoğdu O ym. The effects of therapeutic exercises on pain, muscle strength, functional capacity, balance and hemodynamic parameters in knee osteoarthritis patients: a randomized

controlled study of supervised versus home exercises. *Rheumatol Int.* 2017;37:399-407.

Lefevre-Colau MM, Nguyen C, Haddad R ym. Is physical activity, practiced as recommended for health benefit, a risk factor for osteoarthritis? *Ann Ohys Rehabil Med.* 2016;59:196-206.

Leung YY, Ang LW, Thumboo J, Wang R, Yuan JM, Koh WP. Cigarette smoking and risk of total knee replacement for severe osteoarthritis among Chinese in Singapore--the Singapore Chinese health study. *Osteoarthritis Cartilage.* 2014;22:764-70.

Liu SH, Driban JB, Eaton CB, McAlindon TE, Harrold RL, Lapane KL. Objectively measured physical activity and symptoms change in knee osteoarthritis. *Am J Med.* 2016;129:497-505.

Loew L, Brosseau L, Wells GA ym. Ottawa panel evidence-based clinical practice guidelines for aerobic walking programs in the management of osteoarthritis. *Arch Phys Med Rehabil.* 2012;93:1269-85.

Losina E, Klara K, Michl GL, Collins JE, Katz JN. Development and feasibility of a personalized, interactive risk calculator for knee osteoarthritis. *BMC Musculoskelet Disord.* 2015;16:312.

Lu B, Ahmad O, Zhang FF ym. Soft drink intake and progression of radiographic knee osteoarthritis: data from the osteoarthritis initiative. *BMJ Open.* 2013;3. pii: e002993

Lu M, Su Y, Zhang Z ym. Effectiveness of aquatic exercise for treatment of knee osteoarthritis: systematic review and meta-analysis. *Z Rheumatol.* 2015;74:543-52.

Lu B, Driban JB, Duryea J, McAlindon T, Lapane KL, Eaton CB. Milk consumption and progression of medial tibiofemoral knee osteoarthritis: data from the Osteoarthritis Initiative. *Arthritis Care Res.* 2014 ;66:802-9.

Martel-Pelletier J, Roubille C, Abram F ym. First-line analysis of the effects of treatment on progression of structural changes in knee osteoarthritis over 24 months: data from the osteoarthritis initiative progression cohort. *Ann Rheum Dis*. 2015;74:547-56.

Martin KR, Kuh D, Harris TB, Guralnik JM, Coggon D, Wills AK. Body mass index, occupational activity, and leisure-time physical activity: an exploration of risk factors and modifiers for knee osteoarthritis in the 1946 British birth cohort. *BMC Musculoskelet Disord*. 2013;14:219.

McAlindon TE, Bannuru RR, Sullivan MC ym. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage*. 2014;22:363e388.

Messier SP, Mihalko SL, Legault C ym. Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial. *JAMA*. 2013;310:1263-73.

Mezhov V, Ciccutini FM, Hanna FS, Brennan SL, Wang YY, Urquhart DM, Wluka AE. Does obesity affect knee cartilage? A systematic review of magnetic resonance imaging data. *Obes Rev*. 2014;15:143-57.

Michaëlsson K, Byberg L, Ahlbom A, Melhus H, Farahmand BY. Risk of severe knee and hip osteoarthritis in relation to level of physical exercise: a prospective cohort study of long-distance skiers in Sweden. *PLoS One*. 2011;6:e18339.

Minshull C, Gleeson N. Consideration of the principles of resistance training in exercise studies for the management of knee osteoarthritis; a systematic review. *Ann Phys Med Rehabil*. 2017;Mar 30. pii:S00003-9993(17)30214-9.

Mork PJ, Holtermann A, Nilsen TI. Effect of body mass index and physical exercise on risk of knee and hip osteoarthritis: longitudinal data

from the Norwegian HUNT Study. *J Epidemiol Community Health*. 2012 ;66:678-83.

Multanen J. Exercise for bone and cartilage in postmenopausal women with mild knee osteoarthritis” (Liikuntaa luustolle ja rustoille vaihdevuosi- iän ohittaneilla naisilla, joilla on lievä polven nivelrikko). Väitöskirja 2016. <http://urn.fi/URN:ISBN:978-951-39-6564-8>. Viitattu 23.5.2017.

Muthuri SG, Zhang W, Maciewicz RA, Muir K, Doherty M. Beer and wine consumption and risk of knee or hip osteoarthritis: a case control study. *Arthritis Res Ther*. 2015;17:23.

Nicholls E, Thomas E, van der Windt DA, Croft PR, Peat G. Pain trajectory groups in persons with, or at high risk of, knee osteoarthritis: findings from the Knee Clinical Assessment Study and the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2014;22:2041-50.

Pinto D, Robertson MC, Hansen P, Abbott JH. Cost-effectiveness of nonpharmacologic, nonsurgical interventions for hip and/or knee osteoarthritis: systematic review. *Value Health* . 2012;15:1-12.

Pisters MF, Veenhof C, van Dijk GM, Dekker J; CARPA Study Group. Avoidance of activity and limitations in activities in patients with osteoarthritis of the hip or knee: a 5 year follow-up study on the mediating role of reduced muscle strength. *Osteoarthritis Cartilage*. 2014;22:171-7.

Polven ja lonkan nivelrikon fysioterapia. Hyvä fysioterapiakäytäntö 13.2.2013. Suomen Fysioterapeutit – Finlands Fysioterapeuter ry:n asettama työryhmä Jyrki Kettunen, Petri Salo, Mika Ulaska, Heli Kangas ja Sirpa Ahtola. www.suomenfysioterapeutit.fi. Viitattu 12.11.2017.

Polvi- ja lonkkanivelrikko. Käypä hoito -suositus. 25.08.2014. Suomalaisen Lääkäriseuran Duodecimin ja Suomen Ortopediayhdistys ry:n asettama työryhmä. www.kaypahoito.fi/kh/kaypahoito?suositus=hoi50054. Viitattu 12.11.2017.

Rahmann AE. Exercise for people with hip or knee osteoarthritis: a comparison of land-based and aquatic interventions. *J Sports Med.* 2010;1:123–135.

Reginster JY, Dudler J, Blicharski T, Pavelka K. Pharmaceutical-grade Chondroitin sulfate is as effective as celecoxib and superior to placebo in symptomatic knee osteoarthritis: the ChONDroitin versus CElecoxib versus Placebo Trial (CONCEPT). *Ann Rheum Dis.* 2017;76:1537-1543.

Regnaux JP, Lefevre-Coalu MM, Tringuart L ym. High-intensity versus low-intensity physical activity or exercise in people with hip or knee osteoarthritis. *Cochrane Database Syst. Rev.* 2015;10::CD010203.

Reyes C, Leyland KM, Peat G, Cooper C, Arden NK, Prieto-Alhambra D. Association between overweight and obesity and risk of clinically diagnosed knee, hip, and hand osteoarthritis: A population-based cohort study. *Arthritis Rheumatol.* 2016;68:1869-75.

Richmond SA, Fukuchi RK, Ezzat A, Schneider K, Schneider G, Emery CA. Are joint injury, sport activity, physical activity, obesity, or occupational activities predictors for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther.* 2013;43:515-19.

Runhaar J, van Middelkoop M, Reijman M. ym. Prevention of knee osteoarthritis in overweight females: the first preventive randomized controlled trial in osteoarthritis. *Am J Med.* 2015;128:888-95.

Runhaar J, Deroisy R, van Middelkoop M ym. The role of diet and exercise and glucosamine sulfate in the prevention of knee osteoarthritis: Further results from the Prevention of knee Osteoarthritis in Overweight Females (PROOF) study. *Semin Arthritis Rheum.* 2016;45(4 Suppl):S42-8.

Runhaar J, de Vos BC, van Middelkoop M ym. Prevention of incident knee osteoarthritis by moderate weight loss in overweight and obese females. *Arthritis Care Res.* 2016;68: 1428-33.

Sharma M. Yoga as an alternative and complementary approach for arthritis: a systematic review. *J Evid Based Complementary Altern Med*. 2014;19:51-8.

Sharpe PA, Wilcox S, Schoffman DE, Hutto B4, Ortaglia A. Association of complementary and alternative medicine use with symptoms and physical functional performance among adults with arthritis. *Disabil Health J*. 2016;9:37-45.

Shengelia R, Parker SJ, Ballin M, George T, Reid MC. Complementary therapies for osteoarthritis: are they effective? *Pain Manag Nurs*. 2013;14:e274-88.

Silverwood V, Blagojevic-Bucknall M, Jinks C, Jordan JL, Protheroe J, Jordan KP. Current evidence on risk factors for knee osteoarthritis in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage*. 2015;23:507-15.

Singh JA, Noorbaloochi S, MacDonald R, Maxwell LJ. Chondroitin for osteoarthritis. *Cochrane Database Syst Rev*. 2015;1:CD005614.

Smith BE, Hendrick P3 Smith TO. Should exercises be painful in the management of chronic musculoskeletal pain? A systematic review and meta-analysis. *Br J Sports Med*. 2017;51:1679-1687.

Spahn G, Grosser V, Schiltenswolf M, Schröter F, Grifka J. Football as risk factor for a non-injury-related knee osteoarthritis - results from a systematic review and metaanalysis (in German). *Sportverletz Sportschaden*. 2015;29:27-39.

Teichtahl AJ, Wluka AE, Wang Y ym. Effect of long-term vigorous physical activity on healthy adult knee cartilage. *Med Sci Sports Exerc*. 2012;44:985-92.

Teichtahl AJ, Wluka AE, Tanamas SK, Wang Y, Strauss BJ, Proietto J, Dixon JB, Jones G, Forbes A, Cicuttini FM. Weight change and change in

tibial cartilage volume and symptoms in obese adults. *Ann Rheum Dis.* 2015;74:1024-9.

Timmins KA, Leech RD, Batt ME, Edwards KL. Running and osteoarthritis: a systematic review and meta-analysis. *Am J Sports Med.* 2017;45:1447-1457.

Tran G, Smith TO, Grice A, Kingsbury SR, McCrory P, Conaghan PG. Does sports participation (including level of performance and previous injury) increase risk of osteoarthritis? A systematic review and meta-analysis. *Br J Sports Med.* 2016;50:1459-66.

Zhou ZY, Liu YK, Chen HL, Liu F. Body mass index and knee osteoarthritis risk: a dose-response meta-analysis. *Obesity.* 2014;22:2180-5.

Urquhart DM, Tobing JE, Hanna FS ym. What is the effect of physical activity on the knee joint? A systematic review. *Med Sci Sports Exerc.* 2011;43:432-42.

Uthman OA, van der Windt DA, Jordan JL ym. Exercise for lower limb osteoarthritis: systematic review incorporating trial sequential analysis and network meta-analysis. *BMJ.* 2013;347:f5555

Uusi-Rasi K, Patil R, Karinkanta S, Tokola K, Kannus P, Sievänen H. Exercise training in treatment and rehabilitation of hip osteoarthritis: A 12-week pilot trial. *J Osteoporos.* 2017;2017:3905492.

van der Esch M, Holla JF, van der Leeden M ym. Decrease of muscle strength is associated with increase of activity limitations in early knee osteoarthritis: 3-year results from the cohort hip and cohort knee study. *Arch Phys Med Rehabil.* 2014;95:1962-8.

Vigdorichik MJ, Nepple JJ, Eftekhary R, Leuing M, Clohisy JC. What is the association of elite sporting activities with development of hip osteoarthritis? *Am J Sports Med.* 2017;45:961-964.

Wallace IJ, Worthington S, Felson DT ym. Knee osteoarthritis has doubled in prevalence since the mid-20th century. *Proc Natl Acad Sci U S A*. 2017;114:9332-9336.

Waller B, Munukka M, Rantalainen T ym. Effects of high-intensity resistance aquatic training on body composition and walking speed in women with mild knee osteoarthritis: a 4-month RCT with 12-month follow-up. *Osteoarthritis Cartilage* 2017;Mar 2. pii: S1063-4584(17)30869-5.

Walsh NE, Pearson J, Healey EL. Physiotherapy management of lower limb osteoarthritis. *Br Med Bull*. 2017;122:151-161.

Welling M. Etiological pathways of osteoarthritis. Doctoral Thesis, University of Oulu 2017.

Wesseling J, Bastick AN, ten Wolde S, Kloppenburg M, Lafeber FP, Bierma-Zeinstra SM, Bijlsma JW. Identifying trajectories of pain severity in early symptomatic knee osteoarthritis: A 5-year followup of the Cohort Hip and Cohort Knee (CHECK) Study. *J Rheumatol*. 2015;42:1470-7

White DK, Tudor-Locke C, Zhang Y ym. Daily walking and the risk of incident functional limitation in knee OA: an observational study. *Arthritis Care Res*. 2014;66:1328–36.

White DK, Lee J, Song J, Chang RW, Dunlop D. Potential functional benefit from light intensity physical activity in knee osteoarthritis. *Am J Prev Med*. 2017;53:689-696.

Wilcox S, McClenaghan B, Sharpe PA, Baruth M, Hootman JM, Leith K, Dowda M. The steps to health randomized trial for arthritis: a self-directed exercise versus nutrition control program. *Am J Prev Med*. 2015;48:1-12.

Williamson W, Kluzek S, Roberts N ym. Behavioural physical activity interventions in participants with lower-limb osteoarthritis: a systematic review with meta-analysis *BMJ Open*. 2015;5:e007642.

Wood LR, Blagojevic-Bucknall M, Stynes S ym. Impairment-targeted exercises for older adults with knee pain: a proof-of-principle study (TargET-Knee-Pain). *BMC Musculoskelet Disord.* 2016;17:47.

Øiestad BE, Juhl CB, Eitzen I, Thorlund JB. Knee extensor muscle weakness is a risk factor for development of knee osteoarthritis. A systematic review and meta-analysis. *Osteoarthritis Cartilage.* 2015;23:171-7.

Osteoporoosi eli luukato (ks. myös Kaatumiset ja tasapaino ja Lonkkamurtumat)

Armstrong ME, Cairns BJ, Banks E ym. Different effects of age, adiposity and physical activity on the risk of ankle, wrist and hip fractures in postmenopausal women. *Bone.* 2012;50:1394-400.

Babatunde OO, Forsyth JJ, Gidlow CJ. A meta-analysis of brief high-impact exercises for enhancing bone health in premenopausal women. *Osteoporos Int.* 2012;23:109-19.

Beavers KM, Beavers DP, Martin SB ym. Change in bone mineral density during weight loss with resistance versus aerobic exercise training in older adults. *J Gerontol A Biol Sci Med Sci.* 2017 ;72:1582-1585.

Black DM, Rosen CJ. Postmenopausal osteoporosis. *N Engl J Med.* 2016;374:254-62.

Deere K, Sayers A, Rittweger J, Tobias JH. Habitual levels of high, but not moderate or low, impact activity are positively related to hip BMD and geometry: results from a population-based study of adolescents. *J Bone Miner Res.* 2012;27:1887-95.

Fenton JJ, Robbins JA, Amarnath AL, Franks P. Osteoporosis overtreatment in a regional health care system. *JAMA Intern Med.* 2016;176:391-3.

Ferrucci L, Baroni M, Ranchelli A, Lauretani F, Maggio M, Mecocci P, Ruggiero C. Interaction between bone and muscle in older persons with mobility limitations. *Curr Pharm Des.* 2014;20:3178-97.

Geusens PP, van den Bergh JP. Osteoporosis and osteoarthritis: shared mechanisms and epidemiology. *Curr Opin Rheumatol.* 2016;28:97-103.

Giangregorio LM, Macintyre NJ, Thabane L, Skidmore CJ, Papaioannou A. Exercise for improving outcomes after osteoporotic vertebral fracture. *Cochrane Database Syst Rev.* 2013;1:CD008618.

Giangregorio LM, McGill S, Wark JD ym. Too Fit To Fracture: outcomes of a Delphi consensus process on physical activity and exercise recommendations for adults with osteoporosis with or without vertebral fractures. *Osteoporos Int.* 2015;26:891-910.

Gomez-Bruton A, Montero-Marín J, González-Agüero A ym. The effect of swimming during childhood and adolescence on bone mineral density: a systematic review and meta-analysis. *Sports Med.* 2016;46:365-79.

Gomez-Bruton A, Montero-Marín J, González-Agüero A ym. Swimming and peak bone mineral density: a systematic review and meta-analysis. *J Sports Sci.* 2017 Apr 10:1-13.

Gómez-Cabello A, Ara I, González-Agüero A ym. Effects of training on bone mass in older adults: a systematic review. *Sports Med.* 2012;42:301-25.

Harding AT, Beck BR. Exercise, osteoporosis, and bone geometry. *Sports.* 2017;5:29.

Harper C, Pattinson AL, Fernando HA, Zibellini J, Seimon RV, Sainsbury A. Effects of obesity treatments on bone mineral density, bone turnover and fracture risk in adults with overweight or obesity. *Horm Mol Biol Clin Investig.* 2016;28:133-149.

Harvey NC, Biver E, Kaufman JM ym. The role of calcium supplementation in healthy musculoskeletal ageing : An expert consensus meeting of the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO) and the International Foundation for Osteoporosis (IOF). *Osteoporos Int.* 2017;28:447-462.

Hill TR, Aspray TJ. The role of vitamin D in maintaining bone health in older people. *Ther Adv Musculoskelet Dis.* 2017;9:89-95.

Hinton PS, Rector RS, Linden MA ym. Weight-loss-associated changes in bone mineral density and bone turnover after partial weight regain with or without aerobic exercise in obese women. *Eur J Clin Nutr.* 2012;66:606-12.

Hinton PS, Nigh P, Thyfault J. Effectiveness of resistance training or jumping-exercise to increase bone mineral density in men with low bone mass: A 12-month randomized, clinical trial. *Bone.* 2015;79:203-12.

Howe TE, Shea B, Dawson LJ ym. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database Syst Rev.* 2011;7:CD000333.

Ireland A, J Rittweger J. Exercise for osteoporosis: how to navigate between overeagerness and defeatism. *J Musculoskelet Neuronal Interact.* 2017;17:155-161.

Krishnan C, Choksi P, Peterson MD. Abdominal adiposity and low physical activity are independently and inversely associated with bone mineral density. *Obes Res Clin Pract.* 2017;11:740-746.

Levis S, Gómez-Marín O. Vitamin D and physical function in sedentary older men. *J Am Geriatr Soc.* 2017;65:323-331.

Kelley GA, Kelley KS, Kohrt WM. Effects of ground and joint reaction force exercise on lumbar spine and femoral neck bone mineral density in

postmenopausal women: a meta-analysis of randomized controlled trials. *BMC Musculoskelet Disord*. 2012;13:177.

Kerr C, Bottomley C, Shingler S ym. The importance of physical function to people with osteoporosis. *Osteoporos Int*. 2017;28:1597-1607.

Kohrt WM, Barry DW, Schwartz RS. Muscle forces or gravity: what predominates mechanical loading on bone? *Med Sci Sports Exerc*. 2009;41:2050-55.

Ma D, Wu L, He Z. Effects of walking on the preservation of bone mineral density in perimenopausal and postmenopausal women: a systematic review and meta-analysis. *Menopause*. 2013;20:1216-26.

Marques EA, Mota J, Carvalho J. Exercise effects on bone mineral density in older adults: a meta-analysis of randomized controlled trials. *Age*. 2011;34:1493–1515.

Martyn-St James M, Carroll S. Meta-analysis of walking for preservation of bone mineral density in postmenopausal women. *Bone*. 2008;43:521-31.

Martyn-St James M, Carroll S. A meta-analysis of impact exercise on postmenopausal bone loss: the case for mixed loading exercise programmes. *Br J Sports Med*. 2009;43:898-908.

Martyn-St James M, Carroll S. Effects of different impact exercise modalities on bone mineral density in premenopausal women: a meta-analysis. *J Bone Miner Metab*. 2010;28:251-67.

Matikainen N. Osteoporoosi. *Duodecim* 2016;132:1051-8.

McMillan LB, Zengin A, Ebeling PR, Scott D. Prescribing physical activity for the prevention and treatment of osteoporosis in older adults. *Healthcare*. 2017;5. pii: E85. doi: 10.3390/healthcare5040085.

Morseth B, Emaus N, Wilsgaard T ym. Leisure time physical activity in adulthood is positively associated with bone mineral density 22 years later. The Tromsø study. *Eur J Epidemiol.* 2010;25:325-31.

Morseth B, Ahmed LA, Bjørnerem Å ym. Leisure time physical activity and risk of non-vertebral fracture in men and women aged 55 years and older: the Tromsø Study. *Eur J Epidemiol.* 2012;27:463-71.

Multanen J, Nieminen MT, Häkkinen A ym. Effects of high-impact training on bone and articular cartilage: 12-month randomized controlled quantitative MRI study. *J Bone Miner Res.* 2014;29:192-201.

Multanen J, Rantalainen T, Kautiainen H ym. Effect of progressive high-impact exercise on femoral neck structural strength in postmenopausal women with mild knee osteoarthritis: a 12-month RCT. *Osteoporos Int.* 2017;28:1323-1333.

Määttä M, Terho E, Jokinen H ym. Lifestyle factors and site-specific risk of hip fracture in community dwelling older women--a 13-year prospective population-based cohort study. *BMC Musculoskelet Disord.* 2012;13:173.

Nikander R, Kannus P, Dastidar P ym. Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. *BMC Med.* 2010;8:47.

Olmedillas H, González-Agüero A, Moreno LA, Casajus JA, Vicente-Rodríguez G. Cycling and bone health: a systematic review. *BMC Med.* 2012;10:168.

Osteoporoosi. Käypä hoito -suositus. 24.04.2014. Suomalaisen Lääkäriseuran Duodecimin, Suomen Endokrinologiyhdistyksen ja Suomen Gynekologiyhdistyksen asettama työryhmä. <http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi24065>. Viitattu 24.11.2017.

Oura P, Paananen M, Niinimäki J ym. High-impact exercise in adulthood and vertebral dimensions in midlife - the Northern Finland Birth Cohort 1966 study. *BMC Musculoskelet Disord*. 2017;18:433.

Palermo A, Tuccinardi D, Defeudis G ym. BMI and BMD: The potential interplay between obesity and bone fragility. *Int J Environ Res Public Health*. 2016;13. pii: E544.

Polidoulis I, Beyene J, Cheung AM. The effect of exercise on pQCT parameters of bone structure and strength in postmenopausal women--a systematic review and meta-analysis of randomized controlled trials. *Osteoporos Int*. 2012;23:39-51.

Rauma PH, Koivumaa-Honkanen H, Williams LJ, Tuppurainen MT, Kröger HP, Honkanen RJ. Life satisfaction and bone mineral density among postmenopausal women: cross-sectional and longitudinal associations. *Psychosom Med*. 2014;76:709-15.

Rikkonen T. Physical activity and muscle strength as predictors of bone loss and fractures in perimenopausal women. *Dissertations in Health Sciences* 146. Itä-Suomen yliopisto 2012.

Rizzoli R, Stevenson JC, Bauer JM ym. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: a consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas*. 2014;79:122-32.

Robling AG. Is bone's response to mechanical signals dominated by muscle forces? *Med Sci Sports Exerc*. 2009;41:2044-9.

Saarelainen J. Obesity and bone mineral measurements in postmenopausal women. *Publications of the University of Eastern Finland. Dissertations in Health Sciences* 250. Itä-Suomen yliopisto 2014..

Senter C, Appelle N, Behera SK. Prescribing exercise for women. *Curr Rev Musculoskelet Med*. 2013 ;6:164-72.

Shanbhogue VV, Mitchell DM, Rosen CJ, Bouxsein ML. Type 2 diabetes and the skeleton: new insights into sweet bones. *Lancet Diabetes Endocrinol.* 2016;4:159-73.

Shapses SA, Sukumar D. Bone metabolism in obesity and weight loss. *Annu Rev Nutr.* 2012;32:287–309.

Simas V, Hing W, Pope R, Climstein M. Effects of water-based exercise on bone health of middle-aged and older adults: a systematic review and meta-analysis. *Open Access J Sports Med.* 2017;8:39-60.

Soltani S, Hunter GR, Kazemi A, Shab-Bidar S. The effects of weight loss approaches on bone mineral density in adults: a systematic review and meta-analysis of randomized controlled trials. *Osteoporos Int.* 2016;27:2655-71.

Stokes CS, Lammert F. Vitamin D supplementation: less controversy, more guidance needed. *F1000Res.* 2016 Aug 17;5. pii: F1000 Faculty Rev-2017.

Sun Z, Chen H, Berger MR, Zhang L, Guo H, Huang Y. Effects of tai chi exercise on bone health in perimenopausal and postmenopausal women: a systematic review and meta-analysis. *Osteoporos Int.* 2016;27:2901-11.

Tarantino U, Piccirilli E, Fantini M, Baldi J, Gasbarra E, Bei R. Sarcopenia and fragility fractures: molecular and clinical evidence of the bone-muscle interaction. *J Bone Joint Surg Am.* 2015;97:429-37.

Zhao R, Zhao M, Zhang L. Efficiency of jumping exercise in improving bone mineral density among premenopausal women: a meta-analysis. *Sports Med.* 2014;44:1393-402.

Zhao R, Zhao M, Xu Z. The effects of differing resistance training modes on the preservation of bone mineral density in postmenopausal women: a meta-analysis. *Osteoporos Int.* 2015;26:1605-18.

Zhao R, Xu Z, Zhao M. Effects of oestrogen treatment on skeletal response to exercise in the hips and spine in postmenopausal women: a meta-analysis. *Sports Med.* 2015;45:1163-73.

Zhao R, Zhang M, Zhang Q. The Effectiveness of combined exercise interventions for preventing postmenopausal bone loss: a systematic review and meta-analysis. *J Orthop Sports Phys Ther.* 2017;47:241-251.

Zibellini J, Seimon RV, Lee CM ym. Does diet-induced weight loss lead to bone loss in overweight or obese adults? A systematic review and meta-analysis of clinical trials. *J Bone Miner Res.* 2015;30:2168-78.

Xu J, Lombardi G, Jiao W, Banfi G. Effects of exercise on bone status in female subjects, from young girls to postmenopausal women: an overview of systematic reviews and meta-analyses. *Sports Med.* 2016;46:1165-82.

Upala S, Yong WC, Sanguankeo A. Bone mineral density is decreased in fibromyalgia syndrome: a systematic review and meta-analysis. *Rheumatol Int.* 2017;37:617-622.

Vlachopoulos D, Barker AR, Williams CA ym. The Impact of sport participation on bone mass and geometry in male adolescents. *Med Sci Sports Exerc.* 2017;49:317-326..

Waters DL, Ward AL, Villareal DT. Weight loss in obese adults 65 years and older: a review of the controversy. *Exp Gerontol.* 2013;48:1054-61.

Watson SL, Weeks BK, Weis LJ, Harding AT, Horan SA, Beck BR. High-intensity resistance and impact training improves bone mineral density and physical function in postmenopausal women with osteopenia and osteoporosis: the LIFTMOR randomized controlled trial. *J Bone Miner Res.* 2017 Oct 4. doi: 10.1002/jbmr.3284.

Weiss EP, Jordan RC, Frese EM, Albert SG, Villareal DT. Effects of weight loss on lean mass, strength, bone, and aerobic capacity. *Med Sci Sports Exerc.* 2017;49:206-217.

Whitfield GP, Kohrt WM, Pettee Gabriel KK, Rahbar MH, Kohl HW 3rd. Bone mineral density across a range of physical activity volumes: NHANES 2007-2010. *Med Sci Sports Exerc.* 2015;47:326-34.

Toimintakyky ja toiminnanvajaukset

Artaud F, Singh-Manoux A, Dugravot A, Tzourio C, Elbaz A. Decline in fast gait speed as a predictor of disability in older adults. *J Am Geriatr Soc.* 2015;63:1129-36.

Alexander KP. Walking as a window to risk and resiliency. *Circulation* 2017; 136:644-645.

Alley DE, Shardell MD, Peters KW ym. Grip strength cutpoints for the identification of clinically relevant weakness. *J Gerontol A Biol Sci Med Sci.* 2014;69:559-66.

Artaud F, Dugravot A, Sabia S, Singh-Manoux A, Tzourio C, Elbaz A. Unhealthy behaviours and disability in older adults: three-City Dijon cohort study. *BMJ.* 2013;347:f4240.

Beauchet O, Annweiler C, Callisaya ML ym. Poor gait performance and prediction of dementia: results from a meta-analysis. *J Am Med Dir Assoc.* 2016;17:482-90.

Brach JS, Perera S, Gilmore S ym. Effectiveness of a timing and coordination group exercise program to improve mobility in community-dwelling older adults: a randomized clinical trial. *JAMA Intern Med.* 2017 ;170:1437-44.

Bohm S, Mademli L, Mersmann F, Arampatzis A. predictive and reactive locomotor adaptability in healthy elderly: a systematic review and meta-analysis. *Sports Med.* 2015;45:1759-77.

Brown CJ, Flood KL. Mobility limitation in the older patient. A clinical review. *JAMA* 2013;310:1168-77.

Bruce B, Fries JF, Hubert H. Regular vigorous physical activity and disability development in healthy overweight and normal-weight seniors: a 13-year study. *Am J Public Health* 2008;98:1294-9.

Byrne C, Faure C, Keene DJ, Lamb SE. Ageing, muscle power and physical function: a systematic review and implications for pragmatic training interventions. *Sports Med*. 2016;46:1311-32.

Chakravarty EF, Hubert HB, Krishnan E, Bruce BB, Lingala VB, Fries JF. Lifestyle risk factors predict disability and death in healthy aging adults. *Am J Med*. 2012;125:190-7.

Chmelo EA, Crotts CI, Newman JC ym. Heterogeneity of physical function responses to exercise training in older adults. *J Am Geriatr Soc*. 2015;63:462-9.

Christensen U, Støvring N, Schultz-Larsen K, Schroll M, Avlund K. Functional ability at age 75: is there an impact of physical inactivity from middle age to early old age? *Scand J Med Sci Sports*. 2006;16:245-51.

Cooper AJ, Mishra GD, Kuh D. Physical activity across adulthood and physical performance in midlife : findings from a British birth cohort. *Am J Prev Med*. 2011;41:376-84.

Cooper AJ, Simmons RK, Kuh D, Brage S, Cooper R; NSHD scientific and data collection team. Physical activity, sedentary time and physical capability in early old age: British birth cohort study. *PLoS One*. 2015 ;10:e0126465.

Cooper R, Muniz-Terrera G, Kuh D. Associations of behavioural risk factors and health status with changes in physical capability over 10 years of follow-up: the MRC National Survey of Health and Development. *BMJ Open*. 2016;6:e009962.

Cummings SR, Studenski S, Ferrucci L. A diagnosis of disability – giving mobility clinical viability. *JAMA* 2014;31;2061-3.

de Vries NM, van Ravensberg CD, Hobbelen JS, Olde Rikkert MG, Staal JB, Nijhuis-van der Sanden MW. Effects of physical exercise therapy on mobility, physical functioning, physical activity and quality of life in community-dwelling older adults with impaired mobility, physical disability and/or multi-morbidity: a meta-analysis. *Ageing Res Rev.* 2012;11:136-49.

DiPietro L, Jin Y, Talegawkar S, Matthews CE. The joint associations of sedentary time and physical activity with mobility disability in older people: The NIH-AARP Diet and Health Study. *J Gerontol A Biol Sci Med Sci.* 2017 Aug 30.

Dunlop DD, Song J, Semanik PA ym. Relation of physical activity time to incident disability in community dwelling adults with or at risk of knee arthritis: prospective cohort study. *BMJ.* 2014;348:g2472.

Dunlop DD, Semanik P, Song J, Manheim LM, Shih V, Chang RW. Risk factors for functional decline in older adults with arthritis. *Arthritis Rheum.* 2005;52:1274-82.

Donoghue OA, Savva GM, Cronin H, Kenny RA, Horgan NF. Using timed up and go and usual gait speed to predict incident disability in daily activities among community-dwelling adults aged 65 and older. *Arch Phys Med Rehabil.* 2014;95:1954-61.

Ferrucci L, Cooper R, Shardell M, Simonsick EM, Schrack JA, Kuh D. Age-related change in mobility: perspectives from life course epidemiology and geroscience. *J Gerontol A Biol Sci Med Sci.* 2016 ;71:1184-94.

Fielding RA, Guralnik JM, King AC ym. Dose of physical activity, physical functioning and disability risk in mobility-limited older adults: Results from the LIFE study randomized trial. *PLoS One.* 2017;12:e0182155.

Fimland MS, Vie G, Holtermann A, Krokstad S, Nilsen TIL. Occupational and leisure-time physical activity and risk of disability pension: prospective data from the HUNT Study, Norway. *Occup Environ Med.* 2018;75:23-28.

Foebel AD, Pedersen NL. Genetic influences on functional capacities in aging. *Gerontologist* 2016;56 suppl 2:S18-29.

García-Esquinas E, Andrade E, Martínez-Gómez D, Caballero FF, López-García E, Rodríguez-Artalejo F. Television viewing time as a risk factor for frailty and functional limitations in older adults: results from 2 European prospective cohorts. *Int J Behav Nutr Phys Act.* 2017;14:54.

Gill TM, Gahbauer EA, Murphy TE, Han L, Allore HG. Risk factors and precipitants of long-term disability in community mobility: a cohort study of older persons. *Ann Intern Med.* 2012;156:131-40.

Gill TM, Guralnik JM, Pahor M ym. Effect of structured physical activity on overall burden and transitions between states of major mobility disability in older persons: secondary analysis of a randomized trial. *Ann Intern Med.* 2016;165:833-840.

Hinrichs T, von Bonsdorff MB, Törmäkangas T ym. Inverse effects of midlife occupational and leisure time physical activity on mobility limitation in old age--a 28-year prospective follow-up study. *J Am Geriatr Soc.* 2014;62:812-20.

Hortobágyi T, Lesinski M, Gäbler M, VanSwearingen JM, Malatesta D, Granacher U. Effects of three types of exercise interventions on healthy old adults' gait speed: a systematic review and meta-analysis. *Sports Med.* 2015;45:1627-43.

Hu Y, Pikhart H, Malyutina S. Alcohol consumption and physical functioning among middle-aged and older adults in Central and Eastern Europe: results from the HAPIEE study. *Age Ageing.* 2015;44:84-9.

Jacob ME, Yee LM, Diehr PH ym. Can a healthy lifestyle compress the disabled period in older adults? *J Am Geriatr Soc.* 2016;64:1952-1961.

Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol.* 2014;2:819-29.

Kemmler W, von Stengel S, Engelke K, Häberle L, Mayhew JL, Kalender WA. Exercise, body composition, and functional ability: a randomized controlled trial. *Am J Prev Med.* 2010;38:279-87.

Kikkert LH, Vuillerme N, van Campen JP, Hortobágyi T, Lamoth CJ. Walking ability to predict future cognitive decline in old adults: a scoping review. *Ageing Res Rev.* 2016;27:1-14.

Koivukangas M, Strandberg T, Leskinen R, Keinänen-Kiukaanniemi S, Antikainen R. Vanhuksen gerastenia – tunnista riskipotilas. *Suomen Lääkäril* 2017;72:425-30.

Koster A, Penninx BW, Newman AB ym. Lifestyle factors and incident mobility limitation in obese and non-obese older adults. *Obesity.* 2007;15:3122-32.

Krall JR, Carlson MC, Fried LP, Xue QL. Examining the dynamic, bidirectional associations between cognitive and physical functioning in older adults. *Am J Epidemiol.* 2014;180:838-46.

Kritchevsky SB, de Cabo R. Energy - a hallmark of physical function. *J Gerontol A Biol Sci Med Sci.* 2015 ;70:1333.

LaCroix AZ, Guralnik JM, Berkman LF, Wallace RB, Satterfield S. Maintaining mobility in late life. II. Smoking, alcohol consumption, physical activity, and body mass index. *Am J Epidemiol.* 1993;137:858-69.

Lahti J, Sabia S, Singh-Manoux A ym. Leisure time physical activity and subsequent physical and mental health functioning among midlife Finnish,

- British and Japanese employees: a follow-up study in three occupational cohorts. *BMJ Open*. 2016;6:e009788.
- Lang IA, Guralnik JM, Melzer D. Physical activity in middle-aged adults reduces risks of functional impairment independent of its effect on weight. *J Am Geriatr Soc*. 2007;55:1836-41.
- Lange-Maia BS, Newman AB, Cauley JA ym. Sensorimotor peripheral nerve function and the longitudinal relationship with endurance walking in the Health, Aging and Body Composition Study. *Arch Phys Med Rehabil*. 2016;97:45-52.
- Li Q, Wang S, Milot E, Bergeron P, Ferrucci L, Fried LP, Cohen AA. Homeostatic dysregulation proceeds in parallel in multiple physiological systems. *Aging Cell*. 2015;14:1103-12.
- Liberman K, Forti LN, Beyer I, Bautmans I. The effects of exercise on muscle strength, body composition, physical functioning and the inflammatory profile of older adults: a systematic review. *Curr Opin Clin Nutr Metab Care*. 2017;20:30-53.
- Lin JC, Guerrieri JG, Moore AA. Drinking patterns and the development of functional limitations in older adults: longitudinal analyses of the health and retirement survey. *J Aging Health*. 2011;23:806-21.
- Lindhom V, Lahti J, Rahkonen O, Lahelma E, Lallukka T. Joint association of physical activity and body weight with subsequent physical and mental functioning: a follow-up study. *BMC Public Health*. 2013;13:197.
- Maggio M, Lauretani F, De Vita F ym. Multiple hormonal dysregulation as determinant of low physical performance and mobility in older persons. *Curr Pharm Des*. 2014;20:3119-48.
- Manini TM. Energy expenditure and aging. *Ageing Res Rev*. 2010;9:1-11.

Mankowski RT, Anton SD, Axtell R ym. Device-measured physical activity as a predictor of disability in mobility-limited older adults. *J Am Geriatr Soc.* 2017;65:2251-2256.

McLean RR, Shardell MD, Alley DE ym. Criteria for clinically relevant weakness and low lean mass and their longitudinal association with incident mobility impairment and mortality: the foundation for the National Institutes of Health (FNIH) sarcopenia project. *J Gerontol A Biol Sci Med Sci.* 2014;69:576-83.

Myint PK, Surtees PG, Wainwright NW ym. Modifiable lifestyle behaviors and functional health in the European Prospective Investigation into Cancer (EPIC)-Norfolk population study. *Prev Med.* 2007;44:109-16.

Nelson HD, Nevitt MC, Scott JC, Stone KL, Cummings SR. Smoking, alcohol, and neuromuscular and physical function of older women. Study of Osteoporotic Fractures Research Group. *JAMA.* 1994;272:1825-31.

North TL, Palmer TM, Lewis SJ ym. Effect of smoking on physical and cognitive capability in later life: a multicohort study using observational and genetic approaches. *BMJ Open.* 2015;5:e008393.

Nüesch E, Pablo P, Dale CE ym. Incident disability in older adults: prediction models based on two British prospective cohort studies. *Age Ageing.* 2015;44:275-82.

Pahor M1, Guralnik JM2, Ambrosius WT ym. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA.* 2014;311:2387-96.

Pasanen T, Tolvanen S, Heinonen A, Kujala UM. Exercise therapy for functional capacity in chronic diseases: an overview of meta-analyses of randomised controlled trials. *Br J Sports Med.* 2017;51:1459-1465.

Peeters G, Dobson AJ, Deeg DJ, Brown WJ. A life-course perspective on physical functioning in women. *Bull World Health Organ.* 2013;91:661-70.

Perera S, Patel KV, Rosano C ym. Gait speed predicts incident disability: a pooled analysis. *J Gerontol A Biol Sci Med Sci.* 2016;71:63-71.

Pettee Gabriel K, Sternfeld B, Colvin A, Stewart A, Strotmeyer ES, Cauley JA, Dugan S, Karvonen-Gutierrez C. Physical activity trajectories during midlife and subsequent risk of physical functioning decline in late mid-life: The Study of Women's Health Across the Nation (SWAN). *Prev Med.* 2017;105:287-294.

Pitkälä K. Liikunnan merkitys ikääntyneiden ihmisten toimintakyvylle ja avuntarpeelle. *Suoman Lääkäril.* 2005;60:3865-70.

Pluijm SM, Visser M, Puts MT ym. Unhealthy lifestyles during the life course: association with physical decline in late life. *Aging Clin Exp Res.* 2007;19:75-83.

Rantakokko M, Mänty M, Rantanen T. Mobility decline in old age. *Exerc Sport Sci Rev.* 2013;41:19-25.

Rejeski WJ, IP EH, Bertoni AG ym. Lifestyle change and mobility in obese adults with Type 2 diabetes. *N Engl J Med.* 2012;366:1209-17.

Rosso AL, Studenski SA, Chen WG ym. Aging, the central nervous system, and mobility. *J Gerontol A Biol Sci Med Sci.* 2013;68:1379-86.

Sabia S, Elbaz A, Rouveau N, Brunner EJ, Kivimaki M, Singh-Manoux A. Cumulative associations between midlife health behaviors and physical functioning in early old age: a 17-year prospective cohort study. *J Am Geriatr Soc.* 2014;62:1860-8.

Savela S. Physical activity in midlife and healthrelated quality of life, frailty, telomere length and mortality in old age. *Acta Univewrsitatis Ouluensis D Medica* 1276, 2014. *Acta Universitatis Ouluensis D Medica* 1276. ISBN 978-952-62-0686-8.

Shah RC, Buchman AS, Leurgans S, Boyle PA, Bennett DA. Association of total daily physical activity with disability in community-dwelling older persons: a prospective cohort study. *BMC Geriatr.* 2012;12:63.

Singh MA. Exercise to prevent and treat functional disability. *Clin Geriatr Med.* 2002;18:431-62.

Sjölund BM, Wimo A, Engström M, von Strauss E. Incidence of ADL disability in older persons, physical activities as a protective factor and the need for informal and formal care--results from the SNAC-N Project. *PLoS One.* 2015;10:e0138901.

Schrack JA, Zipunnikov V, Simonsick EM, Studenski S, Ferrucci L. Rising energetic cost of walking predicts gait speed decline with aging. *J Gerontol A Biol Sci Med Sci.* 2016;71:947-53.

Stenholm S, Koster A, Valkeinen H, Patel KV, Bandinelli S, Guralnik JM, Ferrucci L. Association of physical activity history with physical function and mortality in old age. *J Gerontol A Biol Sci Med Sci.* 2016;71:496-501.

Sternfeld B, Colvin A, Stewart A ym. The effect of a healthy lifestyle on future physical functioning in midlife women. *Med Sci Sports Exerc.* 2017;49:274-282.

Strand BH, Mishra G, Kuh D, Guralnik JM, Patel KV. Smoking history and physical performance in midlife: results from the British 1946 birth cohort. *J Gerontol A Biol Sci Med Sci.* 2011;66:142-9.

Strobl R, Müller M, Thorand B ym. Men benefit more from midlife leisure-time physical activity than women regarding the development of late-life disability--results of the KORA-Age study. *Prev Med.* 2014 ;62:8-13.

Tak E, Kuiper R, Chorus A, Hopman-Rock M. Prevention of onset and progression of basic ADL disability by physical activity in community dwelling older adults: a meta-analysis. *Ageing Res Rev.* 2013;12:329-38.

Telakivi T. Toimintakyvyn arvioinnin kehittäminen. Suomen Lääkäril. 2011;66:3127-32.

Tikkanen P, Nykänen I, Lönnroos E, Sipilä S, Sulkava R, Hartikainen S. Physical activity at age of 20-64 years and mobility and muscle strength in old age: a community-based study. J Gerontol A Biol Sci Med Sci. 2012 ;67:905-10.

Toimintakyvyn ulottuvuudet. <https://www.thl.fi/en/tutkimus-ja...ja.../toimintakyky/toimintakyvyn-ulottuvuudet>. Viitattu 15.10.2017.

Tola S. Toimintakyky vai työkyky. Facultas-hanke selventää käsitteitä ja ohjaa käytäntöjä. Duodecim 2008;124:1195-6.

Van Abbema R, De Greef M, Crajé C, Krijnen W, Hobbelen H, Van Der Schans C7. What type, or combination of exercise can improve preferred gait speed in older adults? A meta-analysis. BMC Geriatr. 2015;15:72.

VanSwearingen JM, Studenski SA. Aging, motor skill, and the energy cost of walking: implications for the prevention and treatment of mobility decline in older persons. J Gerontol A Biol Sci Med Sci. 2014;69:1429-36.

Virta L, Lahtela K. Henkilökohtaisen toimintakyvyn arviointimenetelmän soveltuvuus suomalaiseen käytäntöön. Keskeiset havainnot PCA-menetelmästä (Personal Capability Assessment). Sosiaali- ja terveysturvan selosteita 46. Kela 2005. ISBN 951-669-673-2.

Vita AJ, Terry RB, Hubert HB, Fries JF. Aging, health risks, and cumulative disability. N Engl J Med. 1998;338:1035-41.

von Bonsdorf MB. Physical activity as a predictor of disability and social and health service use in older people. Studies in sport, physical education and health. 141 University of Jyväskylä, 2009.

Vuori I. Elintapojen terveystvaikutukset. Duodecim 2015;131:729-36.

Williams ED, Eastwood SV, Tillin T, Hughes AD, Chaturvedi N. The effects of weight and physical activity change over 20 years on later-life objective and self-reported disability. *Int J Epidemiol.* 2014;43:856-65.

Wollesen B, Schulz S, Seydell L, Delbaere K. Does dual task training improve walking performance of older adults with concern of falling? *BMC Geriatr.* 2017;17:213.

Tuki- ja liikuntaelimitysten tapaturmat ja rasitustilat

Bonanno DR, Landorf KB, Munteanu SE, Murley G, Menz HB. Effectiveness of foot orthoses and shock-absorbing insoles for the prevention of injury: a systematic review and meta-analysis. *Br J Sports Med.* 2017;51:86-96.

Hagel BE, Pless IB, Goulet C, Platt RW, Robitaille Y. Effectiveness of helmets in skiers and snowboarders: case-control and case crossover study. *BMJ.* 2005;330:281.

Kallinen M, Markku A. Aging, physical activity and sports injuries. An overview of common sports injuries in the elderly. *Sports Med.* 1995;20:41-52.

Kostiainen A, Haikonen K. (toim.) 2010. Koti – ja vapaa-ajan tapaturmat – Tilastokatsaus 2010. Kotitapaturmien ehkäisykampanja.

<http://www.kotitapaturma.fi/wp-content/uploads/2011/08/Tilastokatsaus-2010-Koti-ja-vapaa-ajan-tapaturmat.pdf>. Viitattu 5.9.2017.

kotitapaturma/2016/03/29164417/Iakkaiden_tarkistuslista1.pdf. Viitattu 10.9.2017.

Kujala U. Rasitusvammat, s. 580-599. Kirjassa Vuori I, Taimela S, Kujala U, toim. Liikuntalääketiede. Kustannus Oy Duodecim 2005, Helsinki.

Liikunta. Käypä hoito -suositus. 13.01.2016. Suomalaisen Lääkäriseura Duodecimin ja Käypä hoito -johtoryhmän asettama työryhmä.

<http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50075> .

Viitattu 16.4.2017.

Little RMD, Paterson DHHumphreys DA, Stathokostas L. A 12-month incidence of exercise-related injuries in previously sedentary community-dwelling older adults following an exercise intervention. *BMJ Open* 2013;3:e002831

Parkkari J. Liikuntatapaturmat, s. 567 – 579. Kirjassa Vuori I, Taimela S, Kujala U. (toim.) Liikuntalääketiede. Kustannus Oy Duodecim 2005, Helsinki.

Parkkari J. Liikunnan turvallisuus s. 236-46. Kirjassa Fogelholm M, Vuori I, Vasankari T, toim. Terveysliikunta. Kustannus Oy Duodecim 2011, Helsinki,.

Parkkari J, Kannus P, Kujala U. Liikuntatapaturmat ja niiden ehkäisy, s. 668-675. Teoksessa Jousimaa J. ym., toim. Lääkärinkäsikirja. 10.p. Kustannus Oy Duodecim 2011, Helsinki.

Persaud N, Coleman E, Zwolakowski D, Lauwers B, Cass D. Nonuse of bicycle helmets and risk of fatal head injury: a proportional mortality, case-control study. *CMAJ*. 2012;184:E921-3.

Punainen Risti. 2011. Ensiapuohjeet.

http://www.redcross.fi/ensiapu/ensiapuohjeet/fi_FI/index/. Luettu 05.09.2017.

Riebe D, Franklin BA, Thompson PD ym. Updating ACSM's recommendations for exercise preparticipation health screening. *Med Sci Sports Exerc*. 2015;47:2473-9.

Smith M, Matheson GO, Meeuwisse WH. Injuries in cross-country skiing: a critical appraisal of the literature. *Sports Med*. 1996;21:239-50.

THL 2013. Iäkkäiden tapaturmat.

http://www.thl.fi/fi_FI/web/pistetapaturmille-fi/iakkaat. Viitattu 5.9.2017.

THL. Tapaturmat Suomessa.

<https://www.thl.fi/fi/web/tapaturmat/tapaturmat-suomessa/yleisimmat-tapaturmatyypit>. Viitattu 10. 9.2017.

van den Bekerom Peter MPJ, Struijs AA, Blankevoort L ym. What Is the evidence for rest, ice, compression, and elevation therapy in the treatment of ankle sprains in adults? *J Athl Train*. 2012; 47: 435–443.

Van Eerd D, Munhall C, Irvin E ym. Effectiveness of workplace interventions in the prevention of upper extremity musculoskeletal disorders and symptoms: an update of the evidence. *Occupational and Environmental Medicine* 2016;73:62-70.

Vriend I , Gouttebauge V, Finch CF, van Mechelen W, Verhagen EALM. Intervention strategies used in sport injury prevention studies: a systematic review identifying studies applying the Haddon matrix. *Sports Med*. 2017;47:2027-2043.

Virtanen K. Pyöräilijän vammat. *Duodecim* 2016;132:1352-6.

Elintavat käyttäytymisenä ja elintapojen muuttamisen tekniikat eli strategiat

Aaltonen S, Rottensteiner M, Kaprio J, Kujala UM. Motives for physical activity among active and inactive persons in their mid-30s. *Scand J Med Sci Sports*. 2014;24:727-35.

Absetz P. Liikkumattomasta liikkujaksi, mutta miten? *Liikunta & Tiede* 2013;50:4-8.

Absetz P, Hankonen N. Elämäntapamuutoksen tukeminen terveydenhuollossa: vaikuttavuus ja keinot. *Duodecim* 2011;127:2265-72.

Absetz P, Hankonen N. Miten auttaa potilaita omaksumaan ja ylläpitämään terveellisiä elämäntapoja. *Duodecim* 2017;133:1015-21.

- Allan JL, McMinn D, Daly M. A bidirectional relationship between executive function and health behavior: evidence, implications, and future directions. *Front Neurosci.* 2016;10:386.
- Allom V, Mullan B, Hagger M. Does inhibitory control training improve health behaviour? A meta-analysis. *Health Psychol Rev.* 2016;10:168-86.
- Artinian NT, Fletcher GF, Mozaffarian D ym. Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association. *Circulation.* 2010;122:406-41.
- Ashford S, Edmunds J, French DP. What is the best way to change self-efficacy to promote lifestyle and recreational physical activity? A systematic review with meta-analysis. *Br J Health Psychol.* 2010;15(Pt 2):265-88.
- Austvoll-Dahlgren A, Lidal IB, Mosdøl A, Odgaard-Jensen J , Denison E, Vist GE. Effects of Information and other communication Initiatives on physical activity with one or more other lifestyle habits targeting the general population. Report from Norwegian Knowledge Centre for the Health Services (NOKC) No. 9-2015.
- Baert V, Gorus E, Mets T, Geerts C, Bautmans I. Motivators and barriers for physical activity in the oldest old: a systematic review. *Ageing Res Rev.* 2011;10:464-74.
- Bauman AE, Reis RS, Sallis JF, Wells JC, Loos RJ, Martin BW; Lancet Physical Activity Series Working Group. Correlates of physical activity: why are some people physically active and others not? *Lancet.* 2012;380(9838):258-71.
- Benn Y, Webb TL, Chang BP, Harkin B. What is the psychological impact of self-weighing? A meta-analysis. *Health Psychol Rev.* 2016;10:187-203.

- Bird EL, Baker G, Mutrie N, Ogilvie D, Sahlqvist S, Powell J. Behavior change techniques used to promote walking and cycling: a systematic review. *Health Psychol.* 2013;32:829-38.
- Boswell RG, Kober H. Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. *Obes Rev.* 2016;17:159-77.
- Buckley J, Cohen JD, Kramer AF, McAuley E, Mullen SP. Cognitive control in the self-regulation of physical activity and sedentary behavior. *Front Hum Neurosci.* 2014;8:747.
- Conklin CA, Soreca I, Kupfer DJ ym. Exercise attenuates negative effects of abstinence during 72 hours of smoking deprivation. *Exp Clin Psychopharmacol.* 2017;25:265-272.
- Cradock KA, ÓLaighin G, Finucane FM, Gainforth HL, Quinlan LR, Ginis KA. Behaviour change techniques targeting both diet and physical activity in type 2 diabetes: A systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2017;14:18.
- Dai S, Wang F, Morrison H. Predictors of decreased physical activity level over time among adults: a longitudinal study. *Am J Prev Med.* 2014;47:123-30.
- de Ridder DT, Lensvelt-Mulders G, Finkenauer C, Stok FM, Baumeister RF. Taking stock of self-control: a meta-analysis of how trait self-control relates to a wide range of behaviors. *Pers Soc Psychol Rev.* 2012;16:76-99.
- Donnachie C, Wyke S, Mutrie N, Hunt K. 'It's like a personal motivator that you carried around wi' you': utilising self-determination theory to understand men's experiences of using pedometers to increase physical activity in a weight management programme. *Int J Behav Nutr Phys Act.* 2017;14:61.
- Duckworth AL, Gendler TS, Gross JJ. Situational strategies for self-control. *Perspect Psychol Sci.* 2016;11:35-55.

Duckworth AL, Seligman MEP. The science and practice of self-control. *Perspect Psychol Sci.* 2017;12:715-8.

Engberg E, Alen M, Kukkonen-Harjula K, Peltonen JE, Tikkanen HO, Pekkarinen H. Life events and change in leisure time physical activity: a systematic review. *Sports Med.* 2012;42:433-47.

Epstein LH, Carr KA, Cavanaugh MD, Paluch RA, Bouton ME. Long-term habituation to food in obese and nonobese women. *Am J Clin Nutr.* 2011;94:371-6.

Epstein LH, Temple JL, Roemmich JN, Bouton ME. Habituation as a determinant of human food intake. *Psychol Rev.* 2009;116:384-407.

Field M, Jones A. Elevated alcohol consumption following alcohol cue exposure is partially mediated by reduced inhibitory control and increased craving. *Psychopharmacology.* 2017;234:2979-2988.

Finkelstein EA, Haaland BA, Bilger M ym. Effectiveness of activity trackers with and without incentives to increase physical activity (TRIPPA): a randomised controlled trial. *Lancet Diabetes Endocrinol.* 2016;4:983-995.

French DP, Cameron E, Benton JS, Deaton C, Harvie M. Can communicating personalised disease risk promote healthy behaviour? A systematic review of systematic reviews. *Ann Behav Med.* 2017;51:718-729

García-Esquinas E, Andrade E, Martínez-Gómez D, Caballero FF, López-García E, Rodríguez-Artalejo F. Television viewing time as a risk factor for frailty and functional limitations in older adults: results from 2 European prospective cohorts. *Int J Behav Nutr Phys Act.* 2017;14:54.

Gardner B. A review and analysis of the use of 'habit' in understanding, predicting and influencing health-related behaviour. *Health Psychol Rev.* 2015;9:277-95.

Gardner B, Lally P. Does intrinsic motivation strengthen physical activity habit? Modeling relationships between self-determination, past behaviour, and habit strength. *J Behav Med.* 2013;36:488-97.

Gardner B, Smith L, Lorencatto F, Hamer M, Biddle SJ. How to reduce sitting time? A review of behaviour change strategies used in sedentary behaviour reduction interventions among adults. *Health Psychol Rev.* 2016;10:89-112.

Gierisch JM, Goode AP, Batch BC ym. The impact of wearable motion sensing technologies on physical activity: a systematic review [Internet]. Washington (DC): Department of Veterans Affairs (US); 2015 Sep.

Goode AP, Hall KS, Batch BC ym. The impact of interventions that integrate accelerometers on physical activity and weight loss: a systematic review. *Ann Behav Med.* 2017;51:79-93.

Greaves CJ, Sheppard KE, Abraham C ym. Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health.* 2011;11:119.

Hagger MS, Chatzisarantis NL. An integrated behavior change model for physical activity. *Exerc Sport Sci Rev.* 2014;42:62-9.

Hall AK, Cole-Lewis H, Bernhardt JM. Mobile text messaging for health: a systematic review of reviews. *Annu Rev Public Health.* 2015;36:393-415.

Hall PA, Fong GT. Temporal self-regulation theory: a neurobiologically informed model for physical activity behavior. *Front Hum Neurosci.* 2015;9:117.

Hardcastle SJ, Hancox J, Hattar A, Maxwell-Smith C, Thøgersen-Ntoumani C, Hagger MS. Motivating the unmotivated: how can health behavior be changed in those unwilling to change? *Front Psychol.* 2015;6:835.

Hollands GJ, Marteau TM, Fletcher PC. Non-conscious processes in changing health-related behaviour: a conceptual analysis and framework. *Health Psychol Rev.* 2016;10:381-394.

Hollands GJ, Shemilt I, Marteau TM ym. Portion, package or tableware size for changing selection and consumption of food, alcohol and tobacco. *Cochrane Database Syst Rev.* 2015;9:CD011045.

Houben K, Jansen A. Training inhibitory control. A recipe for resisting sweet temptations. *Appetite.* 2011;56:345-9.

Horodyska K, Luszczynska A1, van den Berg M ym. Good practice characteristics of diet and physical activity interventions and policies: an umbrella review. *BMC Public Health.* 2015;15:19.

Horodyska K, Luszczynska A, Hayes CB. Implementation conditions for diet and physical activity interventions and policies: an umbrella review. *BMC Public Health.* 2015;15:1250.

Jackson SE, Steptoe A, Wardle J. The influence of partner's behavior on health behavior change: the English Longitudinal Study of Ageing. *JAMA Intern Med.* 2015;175:385-92.

Jones A, Di Lemmaab CG, Robinson E, Christiansen P, Nolan S, Tudur-Smith C, Field M. Inhibitory control training for appetitive behaviour change: A meta-analytic investigation of mechanisms of action and moderators of effectiveness. *Appetite* 2016;97:16-28.

Kaner EF, Beyer FR, Garnett Cym. Personalised digital interventions for reducing hazardous and harmful alcohol consumption in community-dwelling populations. *Cochrane Database Syst Rev.* 2017;9:CD011479.

Keadle SK, Conroy DE, Buman MP, Dunstan DW, Matthews CE. Targeting reductions in sitting time to increase physical activity and improve health. *Med Sci Sports Exerc.* 2017;49:1572-1582.

Keesman M, Aarts H, Häfner M, Papies EK. Mindfulness reduces reactivity to food cues: underlying mechanisms and applications in daily life. *Curr Addict Rep*. 2017;4:151-157.

Kliemann N, Vickerstaff V, Croker H1, Johnson F, Nazareth I, Beeken RJ. The role of self-regulatory skills and automaticity on the effectiveness of a brief weight loss habit-based intervention: secondary analysis of the 10 top tips randomised trial. *Int J Behav Nutr Phys Act*. 2017;14:119.

Kuh D, Karunanathan S, Bergman H, Cooper R. A life-course approach to healthy ageing: maintaining physical capability. *Proc Nutr Soc*. 2014;73:237-48.

Lara J, Evans EH, O'Brien N ym. Association of behaviour change techniques with effectiveness of dietary interventions among adults of retirement age: a systematic review and meta-analysis of randomised controlled trials. *BMC Med*. 2014;12:177.

McDermott MS, Oliver M, Iverson D, Sharma R. Effective techniques for changing physical activity and healthy eating intentions and behaviour: a systematic review and meta-analysis. *Br J Health Psychol*. 2016;21:827-841.

Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Health Psychol*. 2009;28:690-701.

Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci*. 2011;6:42.

Michie S, Wood CE, Johnston M, Abraham C, Francis JJ, Hardeman W. Behaviour change techniques: the development and evaluation of a taxonomic method for reporting and describing behaviour change interventions (a suite of five studies involving consensus methods,

randomised controlled trials and analysis of qualitative data). *Health Technol Assess.* 2015;19:1-188.

Murray JM, Brennan SF, French DP, Patterson CC, Kee F, Hunter RF. Effectiveness of physical activity interventions in achieving behaviour change maintenance in young and middle aged adults: A systematic review and meta-analysis. *Soc Sci Med.* 2017;192:125-133.

O'Brien N, McDonald S, Araújo-Soares V ym. The features of interventions associated with long-term effectiveness of physical activity interventions in adults aged 55-70 years: a systematic review and meta-analysis. *Health Psychol Rev.* 2015;9:417-33.

Olander EK, Fletcher H, Williams S, Atkinson L, Turner A, French DP. What are the most effective techniques in changing obese individuals' physical activity self-efficacy and behaviour: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2013;10:29.

Painter SL, Ahmed R, Hill JO. What matters in weight loss? An in-depth analysis of self-monitoring. *J Med Internet Res.* 2017;19:e160

Papies EK. Health goal priming as a situated intervention tool: how to benefit from nonconscious motivational routes to health behaviour. *Health Psychol Rev.* 2016;10:408-424.

Patja K. Tupakkateollisuuden vanhat keinot uudessa käytössä. *Suomen Lääkäril.* 2017;72:2586-7.

Phillips SM, Cadmus-Bertram L, Rosenberg D, Buman MP, Lynch BM. Wearable technology and physical activity in chronic disease: opportunities and challenges. *Am J Prev Med.* 2018;54:144-150.

Poelman MP, de Vet E, Velema E, Seidell JC, Steenhuis IH. Behavioural strategies to control the amount of food selected and consumed. *Appetite.* 2014;72:156-65.

Poelman MP, de Vet E, Velema E, de Boer MR, Seidell JC, Steenhuis IH. PortionControl@HOME: results of a randomized controlled trial evaluating the effect of a multi-component portion size intervention on portion control behavior and body mass index. *Ann Behav Med*. 2015;49:18-28.

Pope JP, Pelletier L, Guertin C. Starting off on the best foot: A review of message framing and message tailoring, and recommendations for the comprehensive messaging strategy for sustained behavior change. *Health Commun*. 2017;16:1-10.

Rebar AL, Loftus AM, Hagger MS. Cognitive control and the non-conscious regulation of health behavior. *Front Hum Neurosci*. 2015;9:122.

Rhodes RE, Martin AD, Taunton JE, Rhodes EC, Donnelly M, Elliot J. Factors associated with exercise adherence among older adults. An individual perspective. *Sports Med*. 1999;28:397-411.

Rhodes RE, Quinlan A, Mistry CD. Do other goals influence physical activity? A systematic review examining the relationship between other goals and physical activity behavior. *Prev Med*. 2016;91:306-317.

Rolls BJ, Roe LS, James BL, Sanchez CE. Does the incorporation of portion-control strategies in a behavioral program improve weight loss in a 1-year randomized controlled trial? *Int J Obes*. 2017;41:434-442.

Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *Int J Behav Nutr Phys Act*. 2017;14:42.

Schoeppe S, Alley S, Van Lippevelde W ym. Efficacy of interventions that use apps to improve diet, physical activity and sedentary behaviour: a systematic review. *Int J Behav Nutr Phys Act*. 2016;13:127.

Segar M, Taber JM, Patrick H, Thai CL, Oh A. Rethinking physical activity communication: using focus groups to understand women's goals,

- values, and beliefs to improve public health. *BMC Public Health*. 2017;17:462.
- Sheeran P, Maki A, Montanaro E ym. The impact of changing attitudes, norms, and self-efficacy on health-related intentions and behavior: A meta-analysis. *Health Psychol*. 2016;35:1178-1188.
- Siddique J, de Chavez PJ, Craft LL, Freedson P, Spring B. The effect of changes in physical activity on sedentary behavior: results from a randomized lifestyle intervention trial. *Am J Health Promot*. 2017;31:287-295.
- Steenhuis I, Poelman M. Portion size: latest developments and interventions. *Curr Obes Rep*. 2017;6:10-17.
- Stephenson A, McDonough SM, Murphy MH, Nugent CD, Mair JL. Using computer, mobile and wearable technology enhanced interventions to reduce sedentary behaviour: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act*. 2017;14:105.
- St Quinton T, Brunton JA. Implicit processes, self-regulation, and interventions for behavior change. *Front Psychol*. 2017;8:346.
- Teixeira PJ, Carraça EV, Marques MM ym. Successful behavior change in obesity interventions in adults: a systematic review of self-regulation mediators. *BMC Med*. 2015;13:84.
- Zheng Y, Klem ML, Sereika SM, Danford CA, Ewing LJ, Burke LE. Self-weighting in weight management: a systematic literature review. *Obesity*. 2015;23:256-65.
- Van Cappellen P, Rice EL, Catalino LI, Fredrickson BL. Positive affective processes underlie positive health behaviour change. *Psychol Health*. 2018;33:77-97.

Vermeer WM, Steenhuis IH, Poelman MP. Small, medium, large or supersize? The development and evaluation of interventions targeted at portion size. *Int J Obes.* 2014;38 Suppl 1:S13-8.

Wilson K, Senay I, Durantini M. When it comes to lifestyle recommendations, more is sometimes less: a meta-analysis of theoretical assumptions underlying the effectiveness of interventions promoting multiple behavior domain change. *Psychol Bull.* 2015;141:474-509.

Wood W, Runger D. Psychology of habit. *Annu Rev Psychol.* 2016;67:289-314.

.